

(12) **United States Patent**  
**Sasaki et al.**

(10) Patent No.: **US 6,335,018 B1**  
(45) Date of Patent: **\*Jan. 1, 2002**

(54) **HIGH MOLECULAR WEIGHT MAJOR  
OUTER MEMBRANE PROTEIN OF  
MORAXELLA**

(75) Inventors: **Ken Sasaki; Robin E. Harkness;  
Michel H. Klein, all of Willowdale  
(CA)**

(73) Assignee: **Aventis Pasteur Limited, Toronto (CA)**

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **08/431,718**

(22) Filed: **May 1, 1995**

(51) Int. Cl.<sup>7</sup> ..... **C07K 5/00; C07K 14/00;  
A61K 39/02; A61K 39/38**

(52) U.S. Cl. .... **424/251.1; 424/190.1;  
424/251.1; 424/235.1; 424/184.1; 435/172.3;  
530/300; 530/806; 530/825; 530/350; 536/23.5**

(58) Field of Search ..... **424/190.1, 251.1,  
424/235.1, 184.1; 435/172.3; 530/300, 806,  
825, 350; 536/23.5**

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

4,258,029 A 3/1981 Moloney et al.  
4,855,283 A 8/1989 Lockhoff et al.  
5,552,146 A \* 9/1996 Hansen et al.  
5,759,813 A \* 6/1998 Hansen et al.  
5,808,024 A \* 9/1998 Sasaki et al.  
5,981,213 A \* 11/1999 Hansen et al.  
5,993,826 A \* 11/1999 Hansen et al.  
6,214,981 B1 \* 4/2001 Tucker et al.

**FOREIGN PATENT DOCUMENTS**

WO WO A 93 03761 3/1983  
WO WO A 91 09952 7/1991  
WO WO A 93 10214 5/1993

**OTHER PUBLICATIONS**

Bowie et al Science 247:1306-1310, 1990.\*  
Bullen et al.\*  
Samukawa et al, Infection & Immunity 68/3:1569-1573,  
2000.\*  
Aebi, C., et al, "A Protective Epitope of Moraxella Catarrha-  
lis is Encoded by Two Different Genes"—Inf. And Immun.  
Nov. 1997, pp. 4367-4377. vol. 65, No. 11.  
Van Hare, G.F., P.A. Shurin, C.D. Marchant, N.A. Cartelli,  
C.E. Johnson, D. Fulton, S. Carlin, and C.H. Kim. Acute  
otitis media caused by *Branhamella catarrhalis*: biology and  
therapy. Rev. Infect. Dis. 9:16-27.  
Chapman, A.J., D.M. Musher, S. Jonsson, J.E. Clarridge,  
and R.J. Wallace. 1985. Development of bactericidal anti-  
body during *Branhamella catarrhalis* Infection. J. Infect.  
Dis. 151:878-882.

Hager, H., A. Verghese, S. Alvarez, and S.L. Berk. 1987.  
*Branhamella catarrhalis* respiratory infections. Rev. Infect.  
Dis. 9:1140-1149.

McLeod, D.T., F. Ahmad, M.J. Croughan, and M.A. Calder.  
1986. Bronchopulmonary infection due to *M. catarrhalis*.  
Clinical features and therapeutic response. Drugs  
31(Suppl.3):109-112.

Nicotra, B., M. Rivera, J.I. Luman, and R.J. Wallace. 1986.  
*Branhamella catarrhalis* as a lower respiratory tract patho-  
gen in patients with chronic lung disease. Arch.Intern.Med.  
146:890-893.

Ninane, G., J. Joly, and M. Kraytman. 1978. Bronchopul-  
monary infection due to *Branhamella catarrhalis* 11 cases  
assessed by transtracheal puncture. Br.Med.Jr. 1:276-278.

Srinivasan, G., M.J. Raff, W.C. Templeton, S.J. Givens, R.C.  
Graves, and J.C. Mel. 1981. *Branhamella catarrhalis* pneu-  
monia. Report of two cases and review of the literature.  
Am.Rev. Respir. Dis. 123:553-555.

West, M., S.L. Berk, and J.K. Smith. 1982. *Branhamella  
catarrhalis* pneumonia. South.Med.J. 75:1021-1023.

Brorson, J.-E., A. Axelsson, and S.E. Holm. 1976. Studies on  
*Branhamella catarrhalis* (*Neisseria catarrhalis*) with spe-  
cial reference to maxillary sinusitis. Scan. J. Infect. Dis.  
8:151-155.

Evans, F.O., Jr., J.B. Sydnor, W.E.C. Moore, G.R. Moore,  
J.L. Manwaring, A.H. Brill, R.T. Jackson, S. Hanna, J.S.  
Skaar, L.V. Holdeman, G.S. Fitz-Hugh, M.A. Sande, and  
J.M. Gwaltney, Jr. 1975. Sinusitis of the maxillary antrum.  
N.Engl.J.Med. 293:735-739.

Tinkelman, D.G., and H.J. Silk. 1989. Clinical and bacte-  
riologic features of chronic sinusitis in children. Am.J.Dis-  
Child. 143:938-942.

Wald, E.R., C. Byers, N.Guerra, M.Casselbrant, and D.  
Beste. 1989. Subacute sinusitis in children. J.Pediatr.  
115:28-32.

Wald, E.R., G.J. Milmo, A. Bowen, J.Ledesma-Medina, N.  
Salamon, and C.D.Bluestone. 1981. Acute maxillary sinusi-  
tis in children. N.Engl.J.Med. 304:749-754.

Christensen, J.J., and B. Bruun. 1985. Bacteremia caused by  
a beta-lactamase producing strain of *Branhamella catarrha-  
lis*. Acta.Pathol. Microbiol. Immunol. Scand. Sect.B  
93:273-275.

Craig, D.B., and P.A. Wehrle. 1983. *Branhamella catarrha-  
lis* septic arthritis. J. Rheumatol. 10:985-986.

(List continued on next page.)

Primary Examiner—Nita Minnifield

(74) Attorney, Agent, or Firm—Sim & McBurney

(57) **ABSTRACT**

An isolated and purified outer membrane protein of a  
Moraxella strain, particularly *M. catarrhalis*, has a molecu-  
lar mass of about 200 kDa. The about 200 kDa outer  
membrane protein as well as nucleic acid molecules encod-  
ing the same are useful in diagnostic applications and  
immunogenic compositions, particularly for in vivo admin-  
istration to a host to confer protection against disease caused  
by a bacterial pathogen that produces the about 200 kDa  
outer membrane protein or produces a protein capable of  
inducing antibodies in a host specifically reactive with the  
about 200 kDa outer membrane protein.

9 Claims, 17 Drawing Sheets

## OTHER PUBLICATIONS

- Gray, L.D., R.E. Van Scoy, J.P. Anhalt, and P.K.W. Yu. 1989. Wound infection caused by *Branhamella catarrhalis*. J.Clin. Microbiol. 27:818-820.
- Guthrie, R., K. Bakenhaster, R.Nelson, and R. Woskobnick. 1988. *Branhamella catarrhalis* sepsis: a case report and review of the literature. J.Infect.Dis. 158:907-908.
- Hiroshi, S., E.J. Anaissie, N.Khardori, and G.P. Bodey. 1988. *Branhamella catarrhalis* septicemia in patients with leukemia. Cancer 61:2315-2317.
- O'Neill, J.H., and P.W. Mathieson. 1987. Meningitis due to *Branhamella catarrhalis*. Aust. N.Z. J. Med. 17:241-242.
- Murphy, T.F. 1989. The surface of *Branhamella catarrhalis*: a systematic approach to the surface antigens of an emerging pathogen. Pediatr. Infect. Dis. J. 8:S75-S77.
- Klingman, K.L., and T.F. Murphy. 1994. Purification and characterization of a high-molecular-weight outer membrane protein of *Moraxella (Branhamella) catarrhalis*. Infect. Immun. 62:1150-1155.
- Helminen, M.E., I. Maciver, J.L. Latimer, J. Klesney-Tait, L.D. Cope, M. Paris, G.H. McCracken, Jr., and E.J. Hansen. 1994. A large, antigenically conserved protein on the surface of *Moraxella catarrhalis* is a target for protective antibodies. J. Infect. Dis. 170:867-872.
- Panezutti H., O. James, E.J. Hanson, Y. Choi, R.E. Harkness, M.H. Klein and P. Chong, 1993. Identification of surface-exposed B-cell epitopes recognized by *Haemophilus influenzae* type b P1 specific monoclonal antibodies. Infect. Immun. 61: 1867-1872.
- Nixon-George et al. (1990), J. Immunology 144:4798-4802.
- Wiesmuller (1989), Vaccine 8:29-33.
- Deres et al. (1989), Nature 342-561.
- Lockhoff, O. Glycolipids as Immunomodulators: Synthesis and Properties. 1991. Chem. Int. Ed. Engl. 30:1611-1620.
- Journal of Infectious Diseases, 158 (4). 1988. 761-765., XP002013102 Bartos L C et al: "Comparison of the Outer Membrane Proteins of 50 Strains of *Branhamella-Catarrhalis*" see the whole document.
- Science, Apr. 14, 1995, 268 (5208) P221-5, United States, XP002013103 Casey PJ: "Protein lipidation in cell signaling." see the whole document.

\* cited by examiner

Figure 1A






Figure 1B



**Figure 2**


**Figure 3**

Figure 4




FIG. 5.




FIG. 6A

Nucleotide Sequence Between Sali and NcoI

10	CCATGGATAT	20	GGCAGGTGT	30	GCTCGCCTGC	40	CGTATGATGG	50	CGATGACACC	60	CCATTGCCCC	70	CATATCTGTA
80	CGATTGACA	90	TGTGATATGA	100	TTTAAACATGT	110	GACATGATTT	120	AACATTGTTT	130	AATACTGTTG	140	CCATCATTAC
150	CATAAATTAG	160	TAACGCATTT	170	AGTAACGCAT	180	TTGTAAAAAT	190	CATTGCGCCC	200	CTTTATGTGT	210	ATCATATGAA
220	TAGAAATATTA	230	TGATTGTATC	240	TGATTATTGT	250	ATCAGAATGG	260	TGATGCTATA	270	TGATGATGCC	280	TACGAGTTGA
290	TTTGGGTAA	300	TCACTCTATG	310	ATTTGATATA	320	TTTTGAAACT	330	AATCTATTGA	340	CTTAAATCAC	350	CATATGGTTA
360	TAATTTAGCA	370	TAATGGTAGG	380	CTTTTGTAA	390	AAATCACATC	400	GCAATATTGT	410	TCCTACTGTTA	420	CTACCATGCT
430	TGAATGACGA	440	TCCCAATCAC	450	CAGATTCAAT	460	CAAGTGATGT	470	GTTTGTATAC	480	GCACCATTTA	490	CCCTAATTAT
500	TTCAATCAAA	510	TGCCATATGC	520	AGCATGTATC	530	ATTTTTTTAA	540	GGTAAACCAC	550	CATGAATCAC	560	ATCTATAAAG
570	TCATCTTTAA	580	CAAAGCCACA	590	GGCACAATTA	600	TGGCAGTGGC	610	AGAGTACGCC	620	AAATCCCACA	630	GCACGGGGGG



FIG. 6B

640	650	660	670	680	690	700
GGGTAGCTG	TGCTACAGGG	CAAGTTGGCA	GTGTATGCAC	TCTGAGCTTT	GCCCCGATTTG	CCGCGCTCGC
710	720	730	740	750	760	770
TGTCCTCGTG	ATCGGTGCAA	CGCTCAGTGG	CAGTGCTTAT	GCTCAAAAAA	AAGATACCAA	ACATATCGCA
780	790	800	810	820	830	840
ATTGGTGAAC	AAAACAGCC	AAGACGCTCA	GGCACTGCCA	AGCGGACGG	TGATCGAGCC	ATTGCTATTG
850	860	870	880	890	900	910
GTGAAAATGC	TAACGCACAG	GGCGGTCAAG	CCATCGCCAT	CGGTAGTAGT	AATAAAACTG	TCAATGGAAG
920	930	940	950	960	970	980
CAGTTTGGAT	AAGATAGGTA	CCGATGCTAC	GGGTCAAGAG	TCCATCGCCA	TCGGTGGTGA	TGTAAAGGCT
990	1000	1010	1020	1030	1040	1050
AGTGGTGATG	CCTCGATTGC	CATCGGTAGT	GATGACTTAC	ATTTGCTTGA	TCAGCATGGT	AATCCTAAAC
1060	1070	1080	1090	1100	1110	1120
ATCCGAAAGG	TACTCTGATT	AACGATCTTA	TTAACGGCCA	TGCAGTATTA	AAAGAAATAC	GAAGCTCAAA
1130	1140	1150	1160	1170	1180	1190
GGATAATGAT	GTAAAAATATA	GACGCACAAC	CGCAAGCGGA	CACGCCAGTA	CTGCAGTGGG	AGCCATGTCA
1200	1210	1220	1230	1240	1250	1260
TATGCACAGG	GTCATTTTTC	CAACGCCCTTT	GGTACACGGG	CAACAGCTAA	AAGTGCCTAT	TCCTTGGCAG

FIG. 6C

1270	1280	1290	1300	1310	1320	1330
TGGGTCTTGC	CGCCACAGCC	GAGGCCAAT	CTACAATCGC	TATTGGTTCT	GATGCAACAT	CTAGCTCGTT
1340	1350	1360	1370	1380	1390	1400
GGGAGCGATA	GCCCTTGGTG	CAGGTACTCG	TGCTCAGCTA	CAGGGCAGTA	TTGCCCTAGG	TCAAGGTTCT
1410	1420	1430	1440	1450	1460	1470
GTTGTCACTC	AGAGTGATAA	TAATTCTAGA	CCGGCCTATA	CACCAAATAC	CCAGGCACTA	GACCCCAAGT
1480	1490	1500	1510	1520	1530	1540
TTCAAGCCAC	CAATAATACG	AAGCGGGTC	CACTTTCCAT	TGGTAGTAAC	TCTATCAAAC	GTAATAATCAT
1550	1560	1570	1580	1590	1600	1610
CAATGTCGGT	GCAGGTGTTA	ATAAAACCGA	TGCCGGTCAAT	GTGGCACACG	TAGAAGCGGT	GGTGAAGTGG
1620	1630	1640	1650	1660	1670	1680
GCTAAGGAGC	GTAGAAATTAC	TTTTTCAGGGT	GATGATAACA	GTA CTGACGT	AAAAATAGGT	TTGGATAATA
1690	1700	1710	1720	1730	1740	1750
CTTTAACTAT	TAAAGGTGGT	GCAGAGACCA	ACGCATTAA	CGATAATAAT	ATCGGTGTGG	TAAAAGAGGC
1760	1770	1780	1790	1800	1810	1820
TGATAATAGT	GGTCTGAAAG	TTAAACTTGC	TAAAACTTTA	AACAATCTTA	CTGAGGTGAA	TACAACTACA
1830	1840	1850	1860	1870	1880	1890
TTAAATGCCA	CAACCACAGT	TAAAGGTAGGT	AGTAGTAGTA	GTACTACACG	TGAATTATTG	AGTGATAGTT

FIG. 6D

1900	1910	1920	1930	1940	1950	1960
TAACCTTTAC	CCAGCCCAAT	ACAGGCAGTC	AAAGCACAAG	CAAAACCGTC	TATGGCGTTA	ATGGGGTGAA
1970	1980	1990	2000	2010	2020	2030
GTTTACTAAT	AATGCAGAAA	CAACAGCAGC	AATCGGCACT	ACTCGTATTA	CCAGAGATAA	AATTGGCTTT
2040	2050	2060	2070	2080	2090	2100
GCTCGAGATG	GTGATGTTGA	TGAAAAACAA	GCACCATATT	TGGATAAAAA	ACAACTTAAA	GTGGGTAGTG
2110	2120	2130	2140	2150	2160	2170
TTGCAATTAC	CATAGACAAT	GGCATTGATG	CAGGTAATAA	AAAGATCAGT	AATCTTGCCA	AAGGTAGCAG
2180	2190	2200	2210	2220	2230	2240
TGCTAACGAT	GCGGTTACCA	TCGAACAGCT	CAAAGCCGCC	AAGCCTACTT	TAAACGCAGG	CGCTGGCATC
2250	2260	2270	2280	2290	2300	2310
AGTGTCACAC	CTACTGAAAT	ATCAGTTGAT	GCTAAGAGTG	GCAATGTTAC	CGCCCCAACT	TACAAACATTG
2320	2330	2340	2350	2360	2370	2380
GCGTGAAAAC	CACCGAGCTT	AACAGTGATG	GCACTAGTGA	TAAATTTAGT	GTTAAGGGTA	GTGGTACGAA
2390	2400	2410	2420	2430	2440	2450
CAATAGCTTA	GTTACCGCCG	AACATTGGC	AAGCTATCTA	AATGAAGTCA	ATCGAACGGC	TGACAGTGCT
2460	2470	2480	2490	2500	2510	2520
CTACAAAGCT	TTACCGTTAA	AGAAGAAGAC	GATGATGACG	CCAACGCTAT	CACCGTGGCT	AAAGATACGA

FIG. 6E

2530	2540	2550	2560	2570	2580	2590
CAAAAATGC	CGGCGCAGTC	AGCATCTTAA	AACTCAAAGG	TAAAAACGGT	CTAACGGTTG	CTACCAAAAA
2600	2610	2620	2630	2640	2650	2660
AGATGGTACG	GTTACCTTTG	GGCTTAGCCA	AGATAGCGGT	CTGACCATTG	GCAAAAGCAC	CCTAAACAAC
2670	2680	2690	2700	2710	2720	2730
GATGGCTTGA	CTGTTAAAGA	TACCAACGAA	CAAATCCAAG	TCGGTGCTAA	TGGCATTAAA	TTTACTAATG
2740	2750	2760	2770	2780	2790	2800
TGAATGGTAG	TAATCCAGGT	ACTGGCATTG	CAAATACCGC	TCGCATTACC	AGAGATAAAA	TTGGCTTTGC
2810	2820	2830	2840	2850	2860	2870
TGGTCTTGAT	GGTGCAGTTG	ATACAAACAA	ACCTTATCTT	GATCAAGACA	AGCTACAAGT	TGGCAATGTT
2880	2890	2900	2910	2920	2930	2940
AAGATTACCA	ACACTGGCAT	TAACGCAGGT	GGTAAAGCCA	TCACAGGGCT	GTCCCCAACA	CTGCCTAGCA
2950	2960	2970	2980	2990	3000	3010
TTGCCGATCA	AAGTAGCCGC	AACATAGAAC	TGGGCAATAC	AATCCAAGAC	AAAGACAAAT	CCAACGCTGC
3020	3030	3040	3050	3060	3070	3080
CAGCATTAAAT	GATATATTAA	ATACAGGCTT	TAACCTAAAA	AATAATAACA	ACCCCATTTGA	CTTTGTCTCC
3090	3100	3110	3120	3130	3140	3150
ACTTATGACA	TTGTTGACTT	TGCCAATGGC	AATGCCACCA	CCGCCACAGT	AACCCATGAT	ACCGCTAACA

FIG. 6F

3160	3170	3180	3190	3200	3210	3220
AAACCAGTAA	AGTGGTATAT	GATGTGAATG	TGGATGATAC	AACCATTCAT	CTAACAGGCA	CTGATGACAA
3230	3240	3250	3260	3270	3280	3290
TAAAAAACTT	GGCGTCAAAA	CCACCAAACCT	GAACAAAACA	AGTGCCTAATG	GTAATACAGC	AACTAACTTT
3300	3310	3320	3330	3340	3350	3360
AATGTTAACT	CTAGTGATGA	AGATGCCCTT	GTTAACGCCA	AAGACATCGC	CGAAAAATCTA	AACACCCCTAG
3370	3380	3390	3400	3410	3420	3430
CCAAGGAAAT	TCACACCACC	AAAGGCACAG	CAGACACCGC	CCTACAAACC	TTTACCCTTA	AAAAGGTAGA
3440	3450	3460	3470	3480	3490	3500
TGAAAAATAAT	AATGCTGATG	ACGCCAACGC	CATCACCGTG	GGTCAAAAAGA	ACGCAAAATAA	TCAAGTCAAC
3510	3520	3530	3540	3550	3560	3570
ACCCTAACAC	TCAAAGGTGA	AAACGGTCTT	AATATTAAAA	CCGACAAAAA	TGGTACGGTT	ACCTTTGGCA
3580	3590	3600	3610	3620	3630	3640
TTAACACCCAC	AAGCGGTCTT	AAAGCCGGCA	AAAGCACCCCT	AAACGACGGT	GGCTTGCTTA	TTAAAAACCC
3650	3660	3670	3680	3690	3700	3710
CACTGGTAGC	GAACAAATCC	AAGTCGGTGC	TGATGGCGTG	AAGTTTGCCA	AGGTTAATAA	TAATGGTGTT
3720	3730	3740	3750	3760	3770	3780
GTAGGTGCTG	GCATTGATGG	CACAACTCGC	ATTACCAGAG	ATGAAATTGG	CTTTACTGGG	ACTAATGGCT

FIG. 6G

3790	3800	3810	3820	3830	3840	3850
CACTTGATAA	AAGCAAACCC	CACCTAAGCA	AAGACGGCAT	TAACGCAGGT	GGTAAAAGA	TTACCAACAT
3860	3870	3880	3890	3900	3910	3920
TCAATCAGGT	GAGATTGCC	AAAACAGCCA	TGATGCTGTG	ACAGGCGGCA	AGATTATGA	TTTAAAAACC
3930	3940	3950	3960	3970	3980	3990
GAACTTGAAA	ACAAAATCAG	CAGTACTGCC	AAAACAGCAC	AAAACTCATT	ACACGAATTC	TCAGTAGCAG
4000	4010	4020	4030	4040	4050	4060
ATGAACAAGG	TAATAACTTT	ACGGTTAGTA	ACCCTTACTC	CAGTTATGAC	ACCTCAAAGA	CCTCTGATGT
4070	4080	4090	4100	4110	4120	4130
CATCACCTTT	GCAGGTGAAA	ACGGCATTAC	CACCAAGGTA	AATAAAGGTG	TGGTGCGTGT	GGGCATTGAC
4140	4150	4160	4170	4180	4190	4200
CAAAACCAAG	GCTTAACCAC	GCCTAAGCTG	ACCGTGGGTA	ATAATAATGG	CAAAGGCATT	GTCATTGACA
4210	4220	4230	4240	4250	4260	4270
GCCAAAATGG	TCAAAAATACC	ATCACAGGAC	TAAGCAACAC	TCTAGCTAAT	GTTACCAATG	ATAAAGGTAG
4280	4290	4300	4310	4320	4330	4340
CGTACGCACC	ACAGAACAGG	GCAATATAAT	CAAAGACGAA	GACAAAACCC	GTGCCGCCAG	CATTGTTGAT
4350	4360	4370	4380	4390	4400	4410
GTGCTAAGCG	CAGGCTTTAA	CTTGCAAGGC	AATGGTGAAG	CGGTTGACTT	TGCTCTCCACT	TATGACACCG

## FIG. 6H

4420 TCAACTTTGC 4430 CGATGGCAAT 4440 GCCACCACCG 4450 CTAAGGTGAC 4460 CTATGATGAC 4470 ACAAGCAAAA 4480 CCAGTAAAGT

4490 GGTCTATGAT 4500 GTCAATGTGG 4510 ATGATACAAC 4520 CATGAAGTT 4530 AAAGATAAAA 4540 AACTTGGCGT 4550 AAAAACCCACC

4560 ACATTGACCA 4570 GTACTGGCAC 4580 AGGTGCTAAT 4590 AAATTGGCCC 4600 TAAAGCAATCA 4610 AGCTACTGGC 4620 GATGCGCTTG

4630 TCAAGGCCAG 4640 TGATATCGTT 4650 GCTCATCTAA 4660 ACACCTTATC 4670 TGGCGACATC 4680 CAAACTGCCA 4690 AAGGGGCAAG

4700 CCAAGCGAAC 4710 AACTCAGCAG 4720 GCTATGTGGA 4730 TGCTGATGGC 4740 AATAAGGTCA 4750 TCTATGACAG 4760 TACCGATAAC

4770 AAGTACTATC 4780 AAGCCAAAAA 4790 TGATGGCACA 4800 GTTGATAAAA 4810 CCAAAGAAAT 4820 TGCCAAAGAC 4830 AAACCTGGTCG

4840 CCCAAGCCCA 4850 AATCCAGAT 4860 GGCACATTGG 4870 CTCAAATGAA 4880 TGTCAAATCA 4890 GTCATTAACA 4900 AAGAAACAAGT

4910 AAATGATGCC 4920 AATAAAAAGC 4930 AAGGCATCAA 4940 TGAAGACAAC 4950 GCCTTTGTGA 4960 AAGGACTTGA 4970 AAAAGCCGCT

4980 TCTGATAACA 4990 AAACCAAAAA 5000 CGCCGCAGTA 5010 ACTGTGGGTG 5020 ATTTAAATGC 5030 CGTTGCCCAA 5040 ACACCGCTGA

FIG. 6I

5050	5060	5070	5080	5090	5100	5110
CCTTGCAGG	GGATACAGG	ACAACGGCTA	AAAAACTGGG	CGAGACTTTG	ACCATCAAAG	GTGGGCAAAC
5120	5130	5140	5150	5160	5170	5180
AGACACCAAT	AAGCTAACCG	ATAATAACAT	CGGTGTGGTA	GCAGGTACTG	ATGGCTTCAC	TGTCAAACCTT
5190	5200	5210	5220	5230	5240	5250
GCCAAAGACC	TAACCAATCT	TAACAGCGTT	AATGCAGGTG	GCACCAAAAT	TGATGACAAA	GGCGTGTCTT
5260	5270	5280	5290	5300	5310	5320
TTGTGACTC	AAGCGGTCAA	GCCAAAGCAA	ACACCCCTGT	GCTAAGTGCC	AATGGGCTGG	ACCTGGGTGG
5330	5340	5350	5360	5370	5380	5390
CAAGGTCATC	AGTAATGTGG	GCAAAGGCAC	AAAAGATACC	GACGCTGCCA	ATGTACAACA	GTTAAACGAA
5400	5410	5420	5430	5440	5450	5460
GTACGCAACT	TGTTGGGTCT	TGGTAATGCT	GGTAATGATA	ACGCTGACGG	CAATCAGGTA	AACATTGCCG
5470	5480	5490	5500	5510	5520	5530
ACATCAAAA	AGACCCAAAT	TCAGGTTTCAT	CATCTAACCG	CACGTGTCATG	AAAGCAGGCA	CGGTACTTGG
5540	5550	5560	5570	5580	5590	5600
CGGTAAAGGT	AATAACGATA	CCGAAAACCT	TGCCACTGGT	GGTATACAAG	TGGGCGTGGA	TAAAGACGGC
5610	5620	5630	5640	5650	5660	5670
AACGCTAACG	GCGATTTAAG	CAATGTTTGG	GTCAAAAACCC	AAAAAGATGG	CAGCAAAAAA	GCCCTGCTCG
5680	5690	5700	5710	5720	5730	5740
CCACTTATAA	CGCCGCAGGT	CAGACCAACT	ATTTGACCAC	CAACCCCGCA	GAAGCCATTG	ACAGAATAAA




FIG. 6J

5750	5760	5770	5780	5790	5800	5810
TGAACAAGGT	ATCCGCTTCT	TCCATGTCAA	CGATGGCAAT	CAAGAGCCTG	TGGTACAAGG	GCGTAACGGC
5820	5830	5840	5850	5860	5870	5880
ATTGACTCAA	GTGCCCTCAGG	CAAGCACTCA	GTGGCGATAG	GTTTCCAGGC	CAAGGCAGAT	GGTGAAGCCG
5890	5900	5910	5920	5930	5940	5950
CCGTTGCCAT	AGGCAGACAA	ACCCAAGCAG	GCAACCAATC	CATCGCCATC	GGTGATAACG	CACAAGCCAC
5960	5970	5980	5990	6000	6010	6020
GGGCGATCAA	TCCATCGCCA	TCGGTACAGG	CAATGTGGTA	GCAGGTAAGC	ACTCTGGTGC	CATCGGCGAC
6030	6040	6050	6060	6070	6080	6090
CCAAGCACTG	TTAAGGCTGA	TAACAGTTAC	AGTGTGGGTA	ATAACAACCA	GTTTACCGAT	GCCACTCAAA
6100	6110	6120	6130	6140	6150	6160
CCGATGTCTT	TGGTGTGGC	AATAACATCA	CCGTGACCGA	AAGTAACTCG	GTTGCCCTTAG	GTTCAAACTC
6170	6180	6190	6200	6210	6220	6230
TGCCATCAGT	GCAGGCACAC	ACGCAGGCAC	ACAAGCCAAA	AAATCTGACG	GCACAGCAGG	TACAACCACC
6240	6250	6260	6270	6280	6290	6300
ACAGCAGGTG	CAACCGGTAC	GGTTAAAGGC	TTTGCTGGAC	AAACGGCGGT	TGGTGCGGTC	TCCGTGGGTG
6310	6320	6330	6340	6350	6360	6370
CCTCAGGTGC	TGAACGCCGT	ATCCAAAATG	TGGCAGCAGG	TGAGGTCAGT	GCCACCAGCA	CCGATGCGGT
6380	6390	6400	6410	6420	6430	6440
CAATGGTAGC	CAGTTGTACA	AAGCCACCCA	AAGCATTGCC	AACGCAACCA	ATGAGCTTGA	CCATCGTATC

FIG. 6K

6450	6460	6470	6480	6490	6500	6510
CACCAAAACG	AAAATAAGC	CAATGCAGG	ATTTCATCAG	CGATGGCGAT	GGCGTCCATG	CCACAAGCCT
6520	6530	6540	6550	6560	6570	6580
ACATTCCCTGG	CAGATCCATG	GTTACCGGG	GTATTGCCAC	CCACAACGGT	CAAGGTGCGG	TGGCAGTGGG
6590	6600	6610	6620	6630	6640	6650
ACTGTCTGAAG	CTGTCTGGATA	ATGGTCAATG	GGTATTAA	ATCAATGGTT	CAGCCGATAC	CCAAGGCCAT
6660	6670	6680	6690	6700	6710	6720
GTAGGGGCGG	CAGTTGGTGC	AGGTTTTCAC	TTTAAAGCCA	TAAATCGCAA	GATTTACTTT	AAAAATCAAT
6730	6740	6750	6760	6770	6780	6790
CTCACCATAG	TTGTATAAAA	CAGCATCAGC	ATCAGTCATA	TTACTGATGC	TGATGTTTTT	TATCACTTAA
6800	6810	6820	6830	6840	6850	6860
ACCATTTTAC	CGCTCAAGTG	ATTCTCTTTC	ACCATGACCA	AATCGCCATT	GATCATAGGT	AAACTTATTG
6870	6880	6890	6900	6910	6920	6930
AGTAAATTTT	ATCAATGTAG	TTGTTAGATA	TGGTTAAAT	TGTGCCATTG	ACCAAAAAAT	GACCGATTTA
6940	6950	6960	6970			
TCCCCGAAAT	TTCTGATTAT	GATCCGTTGA	CCTGCAGGTC	GAC		

FIG. 7.



1

# HIGH MOLECULAR WEIGHT MAJOR OUTER MEMBRANE PROTEIN OF MORAXELLA

## FIELD OF THE INVENTION

The present invention relates to the field of immunology and is particularly concerned with outer membrane proteins from *Moraxella*, methods of production thereof, genes encoding such proteins and uses thereof.

## BACKGROUND OF THE INVENTION

Otitis media is the most common illness of early childhood with approximately 70% of all children suffering at least one bout of otitis media before the age of seven. Chronic otitis media can lead to hearing, speech and cognitive impairment in children. It is caused by bacterial infection with *Streptococcus pneumoniae* (approximately 50%), non-typable *Haemophilus influenzae* (approximately 30%) and *Moraxella (Branhamella) catarrhalis* (approximately 20%). In the United States alone, treatment of otitis media costs between one and two billion dollars per year for antibiotics and surgical procedures, such as tonsillectomies, adenoidectomies and insertion of tympanostomy tubes. Because otitis media occurs at a time in life when language skills are developing at a rapid pace, developmental disabilities specifically related to learning and auditory perception have been documented in youngsters with frequent otitis media.

*M. catarrhalis* mainly colonizes the respiratory tract and is predominantly a mucosal pathogen. Studies using cultures of middle ear fluid obtained by tympanocentesis have shown that *M. catarrhalis* causes approximately 20% of cases of otitis media (ref. 1—Throughout this application, various references are referred to in parenthesis to more fully describe the state of the art to which this invention pertains. Full bibliographic information for each citation is found at the end of the specification, immediately preceding the claims. The disclosures of these references are hereby incorporated by reference into the present disclosure).

The incidence of otitis media caused by *M. catarrhalis* is increasing. As ways of preventing otitis media caused by pneumococcus and nontypeable *H. influenzae* are developed, the relative importance of *M. catarrhalis* as a cause of otitis media can be expected to further increase.

*M. catarrhalis* is also an important cause of lower respiratory tract infections in adults, particularly in the setting of chronic bronchitis and emphysema (refs. 2, 3, 4, 5, 6, 7, and 8). *M. catarrhalis* also causes sinusitis in children and adults (refs. 9, 10, 11, 12, and 13) and occasionally causes invasive disease (refs. 14, 15, 16, 17, 18, and 19).

Like other Gram-negative bacteria, the outer membrane of *M. catarrhalis* consists of phospholipids, lipopolysaccharide (LPS), and outer membrane proteins (OMPs). Eight of the *M. catarrhalis* OMPs have been identified as major components. These are designated by letters A through H, beginning with OMP A which has a molecular mass of 98 kDa to OMP H which has a molecular mass of 21 kDa (ref. 20).

Recently, a high-molecular-weight outer membrane protein of *M. catarrhalis* was purified and characterized (ref. 21). The apparent molecular mass of this protein varies from 350 kDa to 720 kDa as judged by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). This protein appears to be an oligomer of much smaller proteins or subunits thereof of molecular mass 120 to 140 kDa and is antigenically conserved among strains of *Moraxella*.

2

A similar protein named UspA was also reported to be present on the surface of this species of bacteria with an apparent molecular mass of 300 to 400 kDa (ref. 22). Judging from the molecular mass, these two proteins may be the same.

*M. catarrhalis* infection may lead to serious disease. It would be advantageous to provide other outer membrane proteins for *M. catarrhalis* and genes encoding such proteins for use as antigens in immunogenic preparations including vaccines, carriers for other antigens and immunogens and the generation of diagnostic reagents.

## SUMMARY OF THE INVENTION

The present invention is directed towards the provision of purified and isolated major outer membrane protein of *Moraxella catarrhalis* and other *Moraxella* strains, having an apparent molecular mass of about 200 kDa.

In accordance with one aspect of the invention, there is provided an isolated and purified, outer membrane protein of a *Moraxella* strain having a molecular weight of about 200 kDa, as determined by SDS-PAGE, or a fragment or an analog thereof. The outer membrane protein may be substantially in its native conformation (so as to have substantially retained the characteristic immunogenicity of the outer membrane protein in the *Moraxella* strain) and may be isolated from a *M. catarrhalis* strain, such as from *M. catarrhalis* 4223. Such isolated and purified about 200 kDa outer membrane protein is substantially free from non-200 kDa outer membrane protein, phospholipids and lipopolysaccharide of *Moraxella*. The about 200 kDa outer membrane protein is at least about 70 wt % pure, preferably at least about 90 wt % pure, and may be in the form of an aqueous solution thereof.

The present invention also provides a purified and isolated nucleic acid molecule encoding an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, or a fragment or an analog of the outer membrane protein. The protein encoded by the nucleic acid molecule may encode a protein containing the amino acid sequence NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No: 2) for *Moraxella catarrhalis* strain 4223 or containing the corresponding amino acid sequence from other *Moraxella* strains.

In a further aspect of the present invention, there is provided a purified and isolated nucleic acid molecule having a sequence selected from the group consisting of (a) the sequence set out in FIG. 6 (SEQ ID No: 1), or the complementary sequence thereto; (b) a sequence encoding an about 200 kDa protein of a strain of *Moraxella* and containing the amino acid sequence NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No: 2), or the complementary sequence thereto; and (c) a nucleotide sequence which hybridizes under stringent conditions to any one of the sequences defined in (a) or (b). The nucleic acid preferably defined in (c) has at least about 90% sequence identity with any one of the sequences defined in (a) or (b).

The nucleic acid molecules provided herein may be included in a vector adapted for transformation of a host. The nucleic acid molecules provided herein also may be included in an expression vector adapted for transformation of a host along with expression means operatively coupled to the nucleic acid molecule for expression by the host of the about 200 kDa outer membrane protein of a strain of *Moraxella* or the fragment or the analog of the outer mem-

brane protein. A transformed host containing the expression vector is included within the invention, along with a recombinant outer membrane protein or fragment or analog thereof producible by the transformed host.

The expression means may include a nucleic acid portion encoding a leader sequence for secretion from the host of the outer membrane protein or the fragment or the analog of the outer membrane protein. The expression means may include a nucleic acid portion encoding a lipidation signal for expression from the host of a lipidated form of the outer membrane protein or the fragment or analog thereof.

The present invention further includes a live vector for delivery of the outer membrane protein of the invention or a fragment or analog thereof, comprising a vector containing the nucleic acid molecule provided herein. The live vector may be selected from the group consisting of *E. coli*, *Salmonella*, BCG, adenovirus, poxvirus, vaccinia and poliovirus.

In accordance with a further aspect of the present invention, there is provided a peptide having no less than six amino acids and no more than 150 amino acids and containing an amino acid sequence corresponding to a portion only of the outer membrane protein of the invention, or a fragment or analog thereof. The peptide may be one having the amino acid sequence NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No: 2) for the *Moraxella catarrhalis* 4283 strain or the amino acid sequence for the corresponding peptide for other strains of *Moraxella*.

The present invention also provides an immunogenic composition comprising an immunoeffective amount of an active component, which may be the outer membrane protein or fragment or analog thereof, nucleic acid molecules, recombinant outer membrane proteins, fragments or analogs thereof, live vectors, and/or peptides as provided herein, along with a pharmaceutically acceptable carrier therefor with the active component producing an immune response when administered to a host, which may be a primate, particularly a human. The immunogenic composition may be formulated as a vaccine for in vivo administration to a host to confer protection against diseases caused by a bacterial pathogen that produces the about 200 kDa outer membrane protein or produces a protein capable of inducing antibodies in the host specifically reactive with the about 200 kDa outer membrane protein. In particular, the bacterial pathogen is a strain of *Moraxella*, particularly *M. catarrhalis*. The immunogenic composition may be formulated as a microparticle capsule, ISCOM or liposome preparation. The immunogenic composition may be used in combination with a targeting molecule for delivery to specific cells of the immune system as to mucosal surfaces. The immunogenic compositions of the invention (including vaccines) may further comprise at least one other immunogenic or immunostimulating material and the immunostimulating material may be at least one adjuvant. Suitable adjuvants for use in the present invention include, (but are not limited to) aluminum phosphate, aluminum hydroxide, QS21, Quil A, calcium phosphate, calcium hydroxide, zinc hydroxide, a glycolipid analog, and octadecyl ester of an amino acid, a muramyl dipeptide and a lipoprotein. Advantageous combinations of adjuvants are described in copending U.S. patent application Ser. No. 261,194 filed Jun. 16, 1994, assigned to the assignee hereof and the disclosure of which is incorporated herein by reference. The invention further includes an antibody specific for the outer membrane protein provided herein producible by immunizing a host with an immunogenic composition provided herein.

In a further aspect of the invention, there is provided a method of generating an immune response in a host comprising administering thereto an immuno-effective amount of the immunogenic composition as provided herein. The immune response may be a humoral or a cell-mediated immune response. The immune response may provide protection to the host against diseases caused by a bacterial pathogen that produces the about 200 kDa outer membrane protein or produces a protein capable of inducing antibodies in the host specifically reactive with the about 200 kDa outer membrane protein. Hosts in which protection against disease may be conferred include primates including humans.

The present invention provides, in an additional aspect thereof, a method of producing a vaccine comprising administering the immunogenic composition provided herein to a test host to determine an amount and a frequency of administration of the active component to confer protection against disease caused by a bacterial pathogen that produces the about 200 kDa outer membrane protein or produces a protein capable of inducing antibodies in the host specifically reactive with the about 200 kDa outer membrane protein, and formulating the active component in a form suitable for administration to a treated host in accordance with said determined amount and frequency of administration. The treated host may be a human.

A further aspect of the present invention provides a method of determining the presence of nucleic acid encoding an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, in a sample, comprising the steps of:

- (a) contacting the sample with the nucleic acid molecule provided herein to produce duplexes comprising the nucleic acid molecules and any said nucleic acid molecule encoding the outer membrane protein present in the sample and specifically hybridizable therewith; and
- (b) determining the production of the duplexes.

In yet a further aspect of the invention, there is provided a method of determining the presence of antibodies specifically reactive with outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, in a sample, comprising the steps of:

- (a) contacting the sample with the outer membrane protein as provided herein to produce complexes comprising the outer membrane protein and any said antibodies present in the sample specifically reactive therewith; and
- (b) determining production of the complexes.

In a further aspect of the invention, there is also provided a method of determining the presence of an outer membrane protein if a strain of *Moraxella* having a molecular mass of about 200 kDa, in a sample comprising the steps of:

- (a) immunizing a subject with the immunogenic composition as provided herein, to produce antibodies specific for the outer membrane protein;
- (b) contacting the sample with the antibodies to produce complexes comprising any outer membrane protein present in the sample and said outer membrane protein specific antibodies; and
- (c) determining production of the complexes.

The outer membrane protein may be part of a *Moraxella catarrhalis* strain.

The present invention provides, in a yet further aspect, a diagnostic kit for determining the presence of nucleic acid encoding an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, in a sample, comprising:

5

- (a) the nucleic acid molecule as provided herein;
- (b) means for contacting the nucleic acid with the sample to produce duplexes comprising the nucleic acid molecule and any said nucleic acid present in the sample and hybridizable with the nucleic acid molecule; and
- (c) means for determining production of the duplexes.

In yet a further aspect of the invention, there is provided a diagnostic kit for determining the presence of antibodies in a sample specifically reactive with the outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, comprising:

- (a) the outer membrane protein as provided herein;
- (b) means for contacting the outer membrane protein with the sample to produce complexes comprising the outer membrane protein and any said antibodies present in the sample; and
- (c) means for determining production of the complexes.

The invention also provides a diagnostic kit for detecting the presence of an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, in a sample, comprising:

- (a) an antibody specific for the about 200 kDa outer membrane protein as provided herein;
- (b) means for contacting the antibody with the sample to produce a complex comprising the outer membrane protein and outer membrane-specific antibody; and
- (c) means for determining production of the complex.

In a further aspect of the invention, there is provided a method of producing an isolated and purified outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, is determined by SDS-PAGE, comprising the steps of:

- (a) providing a cell mass of the *Moraxella* strain;
- (b) disrupting the cell mass to provide a cell lysate;
- (c) fractionating the cell lysate to provide a fraction containing the outer membrane protein substantially free from other cell lysate components, and
- (d) recovering said outer membrane protein.

The bacterial strain may be *Moraxella catarrhalis*. The cell lysate may be fractionated by gel electrophoresis.

In this application, the term "about 200 kDa protein" is used to define a family of outer membrane proteins of *M. catarrhalis* having molecular mass of between about 160 and 230 kDa and includes proteins having variations in their amino acid sequences including those naturally occurring in various strains of *Moraxella*. The purified and isolated DNA molecules comprising a gene having an open reading frame of the about 200 kDa protein of the present invention also include those encoding functional analogs of the about 200 kDa protein. In this application, a first protein or peptide is a "functional analog" of a second protein if the first protein is immunologically related to and/or has the same function as the second protein or peptide. The functional analog may be, for example, a fragment of the protein or a substitution, addition or deletion mutant thereof.

Advantages of the present invention include:

- a method for isolating purified about 200 kDa outer membrane protein of a *Moraxella* strain that produces the outer membrane protein, including *Moraxella catarrhalis*;
- an isolated and purified about 200 kDa outer membrane protein isolatable from a *Moraxella* strain; and
- diagnostic kits and immunological reagents for specific identification of *Moraxella* and hosts infected thereby.

6

## BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A and 1B show an analysis of *Moraxella catarrhalis* cell proteins by SDS-PAGE. The identification of the lanes and the sources of the proteins are given in Example 2 below;

FIG. 2 shows a comparative analysis of cell proteins from a number of *M. catarrhalis* strains by SDS-PAGE analysis. The identification of the lanes and the sources of the proteins are given in Example 4 below;

FIG. 3 shows an analysis of isolated and purified about 200 kDa outer membrane protein of *M. catarrhalis* by SDS-PAGE. The identification of the lanes is given in Example 4 below;

FIG. 4 shows the specific recognition of about 200 kDa outer membrane protein by anti-peptide antiserum. The identification of the lanes and antiserum are given in Example 8 below;

FIG. 5 shows restriction maps of clones containing a gene having an open reading frame of the about 200 kDa outer membrane protein of *M. catarrhalis*. An open reading frame of the about 200 kDa outer membrane protein is indicated by the hatched box;

FIGS. 6A to 6K show the nucleotide sequence (SEQ ID No: 1) of the gene having an open reading frame of the about 200 kDa outer membrane protein of *M. catarrhalis*;

FIG. 7 shows the specific identification of *M. catarrhalis* expressing the about 200 kDa outer membrane protein by guinea pig anti-200 kDa specific antiserum in contrast to other bacteria. Identification of the lanes and bacteria appears below.

## GENERAL DESCRIPTION OF INVENTION

Referring to FIG. 1A and 1B and FIG. 2, there is illustrated the separation of a novel outer membrane protein from a variety of strains of *M. catarrhalis* having a molecular mass about 200 kDa. FIG. 3 shows the isolated and purified outer membrane protein.

The purified protein was eluted from the gel and used to raise antibodies in guinea pigs. The antibodies specifically recognize only strains of *M. catarrhalis* which produce the outer membrane protein (Table I below).

Referring to FIG. 4, there is shown the recognition of the about 200 kDa outer membrane protein by antibodies raised in guinea pigs to a synthesized peptide corresponding to an internal fragment of the about 200 kDa protein. The synthesized peptide had the amino acid sequence NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys (SEQ ID No: 3).

Referring to FIG. 5, there is shown restriction maps of clones containing a gene having an open reading frame of the about 200 kDa outer membrane protein. The nucleotide sequence (SEQ ID No: 1) of the gene having an open reading frame coding for the about 200 kDa outer membrane protein as shown in FIG. 6.

In one embodiment of the present invention, the isolated and purified about 200 kDa outer membrane protein as provided herein is useful for generating antibodies that can be used to specifically distinguish *M. catarrhalis* from other bacterial pathogens that cause otitis media and other diseases. Thus referring to FIG. 7, there is illustrated an immunoblot showing the specific reactivity of a guinea pig monospecific anti-200 kDa outer membrane protein antise

rum produced by immunizing mice with the purified about 200 kDa outer membrane protein as provided herein. The bacterial lysates analyzed were as follows:

Lane	Bacterium	Source
1	Molecular Weight Standard	
2	<i>M. catarrhalis</i> 4223	middle ear fluid
3	<i>M. catarrhalis</i> RH408	non-clumping variant of strain 4223
4	<i>H. influenzae</i> , MinnA strain	meningitis isolate
5	non-typable <i>H. influenzae</i> , SB12 strain	otitis media isolate
6	non-typable <i>H. influenzae</i> , SB33 strain	otitis media isolate
7	<i>S. pneumoniae</i> type 6	ATCC 6306
8	<i>S. pneumoniae</i> type J4	ATCC 6314
9	<i>P. aeruginosa</i>	
10	<i>E. coli</i> DH5a	

The results shown in FIG. 7 clearly show the usefulness of outer membrane-specific antisera as provided herein to distinguish between bacterial pathogens that produce diseases with similar clinical symptoms.

Thus, in accordance with another aspect of the present invention, there is provided a vaccine against *Moraxella*, comprising an immunogenically-effective amount of the outer membrane protein as provided herein and a physiologically-acceptable carrier therefor. The outer membrane protein provided herein also may be used as a carrier protein for hapten, polysaccharides or peptides to make a conjugate vaccine against antigenic determinants unrelated to the about 200 kDa outer membrane protein.

The about 200 kDa outer membrane protein provided herein is useful as a diagnostic reagent, as an antigen for the generation of anti-outer membrane protein antibodies, or as an antigen for vaccination against the diseases caused by species of *Moraxella* for detecting infection by *Moraxella*.

In additional embodiments of the present invention, the about 200 kDa outer membrane protein as provided herein may be used as a carrier molecule to prepare chimeric molecules and conjugate vaccines (including glycoconjugates) against pathogenic bacteria, including encapsulated bacteria. Thus, for example, glycoconjugates of the present invention may be used to confer protection against disease and infection caused by any bacteria having polysaccharide antigens including lipooligosaccharides (LOS) and polyribosylphosphate (PRP). Such bacterial pathogens may include, for example, *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Escherichia coli*, *Neisseria meningitidis*, *Salmonella typhi*, *Streptococcus mutants*, *Cryptococcus neoformans*, *Klebsiella*, *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Particular antigens which can be conjugated to outer membrane protein and methods to achieve such conjugations are described in published PCT application WO 94/12641, assigned to the assignee hereof and the disclosure of which is hereby incorporated by reference thereto.

In another embodiment, the carrier function of the outer membrane protein may be used, for example, to induce immunity toward abnormal polysaccharides of tumor cells, or to produce anti-tumor antibodies that can be conjugated to chemotherapeutic or bioactive agents.

It is clearly apparent to one skilled in the art, that the various embodiments of the present invention have many applications in the fields of vaccination, diagnosis, treatment of *Moraxella* infections, and in the generation of immunological reagents. A further non-limiting discussion of such uses is further presented below.

## 1. Vaccine Preparation and Use

Immunogenic compositions, suitable to be used as vaccines, may be prepared from the about 200 kDa outer membrane protein as disclosed herein, which may be purified from the bacteria or which may be produced recombinantly, as well as immunological fragments thereof. The vaccine elicits an immune response in a subject which produces antibodies, including anti-200 kDa outer membrane protein antibodies and antibodies that are opsonizing or bactericidal. Should the vaccinated subject be challenged by *Moraxella* or other bacteria that produce proteins capable of producing antibodies that specifically recognize 200 kDa outer membrane protein, the antibodies bind to and inactivate the bacterium. Furthermore, opsonizing or bactericidal anti-200 kDa outer membrane protein antibodies may also provide protection by alternative mechanisms.

Immunogenic compositions including vaccines may be prepared as injectables, as liquid solutions or emulsions. The about 200 kDa outer membrane protein may be mixed with pharmaceutically acceptable excipients which are compatible with the about 200 kDa outer membrane protein. Such excipients may include, water, saline, dextrose, glycerol, ethanol, and combinations thereof. The immunogenic compositions and vaccines may further contain auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, or adjuvants to enhance the effectiveness thereof. Immunogenic compositions and vaccines may be administered parenterally, by injection subcutaneously or intramuscularly. Alternatively, the immunogenic compositions formed according to the present invention, may be formulated and delivered in a manner to evoke an immune response at mucosal surfaces. Thus, the immunogenic composition may be administered to mucosal surfaces by, for example, the nasal or oral (intra-gastric) routes. Alternatively, other modes of administration including suppositories and oral formulations may be desirable. For suppositories, binders and carriers may include, for example, polyalkylene glycols or triglycerides. Oral formulations may include normally employed excipients such as, for example, pharmaceutical grades of saccharine, cellulose and magnesium carbonate. These compositions can take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 1 to 95% of the about 200 kDa outer membrane protein. The immunogenic preparations and vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective, protective and immunogenic. The quantity to be administered depends on the subject to be treated, including, for example, the capacity of the individual's immune system to synthesize antibodies, and if needed, to produce a cell-mediated immune response. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are readily determinable by one skilled in the art and may be of the order of micrograms of the about 200 kDa outer membrane protein. Suitable regimes for initial administration and booster doses are also variable, but may include an initial administration followed by subsequent administrations. The dosage may also depend on the route of administration and will vary according to the size of the host.

The concentration of the about 200 kDa outer membrane antigen in an immunogenic composition according to the invention is in general about 1 to 95%. A vaccine which contains antigenic material of only one pathogen is a monovalent vaccine. Vaccines which contain antigenic material of several pathogens are combined vaccines and

also belong to the present invention. Such combined vaccines contain, for example, material from various pathogens or from various strains of the same pathogen, or from combinations of various pathogens.

Immunogenicity can be significantly improved if the antigens are co-administered with adjuvants, commonly used as 0.05 to 0.1 percent solution in phosphate-buffered saline. Adjuvants enhance the immunogenicity of an antigen but are not necessarily immunogenic themselves. Adjuvants may act by retaining the antigen locally near the site of administration to produce a depot effect facilitating a slow, sustained release of antigen to cells of the immune system. Adjuvants can also attract cells of the immune system to an antigen depot and stimulate such cells to elicit immune responses.

Immunostimulatory agents or adjuvants have been used for many years to improve the host immune responses to, for example, vaccines. Intrinsic adjuvants, such as lipopolysaccharides, normally are the components of the killed or attenuated bacteria used as vaccines. Extrinsic adjuvants are immunomodulators which are typically non-covalently linked to antigens and are formulated to enhance the host immune responses. Thus, adjuvants have been identified that enhance the immune response to antigens delivered parenterally. Some of these adjuvants are toxic, however, and can cause undesirable side-effects, making them unsuitable for use in humans and many animals. Indeed, only aluminum hydroxide and aluminum phosphate (collectively commonly referred to as alum) are routinely used as adjuvants in human and veterinary vaccines. The efficacy of alum in increasing antibody responses to diphtheria and tetanus toxoids is well established and, more recently, a HBsAg vaccine has been adjuvanted with alum. While the usefulness of alum is well established for some applications, it has limitations. For example, alum is ineffective for influenza vaccination and inconsistently elicits a cell mediated immune response.

A wide range of extrinsic adjuvants can provoke potent immune responses to antigens. These include saponins complexed to membrane protein antigens (immune stimulating complexes), pluronic polymers with mineral oil, killed mycobacteria in mineral oil, Freund's complete adjuvant, bacterial products, such as muramyl dipeptide (MDP) and lipopolysaccharide (LPS), as well as lipid A, and liposomes.

To efficiently induce humoral immune responses (HIR) and cell-mediated immunity (CMI), immunogens are emulsified in adjuvants. Many adjuvants are toxic, inducing granulomas, acute and chronic inflammations (Freund's complete adjuvant) FCA, cytotoxicity (saponins and Pluronic polymers) and pyrogenicity, arthritis and anterior uveitis (LPS and MDP). Although FCA is an excellent adjuvant and widely used in research, it is not licensed for use in human or veterinary vaccines because of its toxicity.

Desirable characteristics of ideal adjuvants include:

- (1) lack of toxicity;
- (2) ability to stimulate a long-lasting immune response;
- (3) simplicity of manufacture and stability in long-term storage;
- (4) ability to elicit both CMI and HIR to antigens administered by various routes, if required;
- (5) synergy with other adjuvants;
- (6) capability of selectively interacting with populations of antigen presenting cells (APC);
- (7) ability to specifically elicit appropriate  $T_H1$  or  $T_H2$  cell-specific immune responses; and
- (8) ability to selectively increase appropriate antibody isotype levels (for example, IgA) against antigens.

U.S. Pat. No. 4,855,283 granted to Lockhoff et al on Aug. 8, 1989 which is incorporated herein by reference thereto, teaches glycolipid analogues including N-glycosylamides, N-glycosylureas and N-glycosylcarbamates, each of which is substituted in the sugar residue by an amino acid, as immuno-modulators or adjuvants. Thus, Lockhoff et al. (U.S. Pat. No. 4,855,283 and ref. 27) reported that N-glycolipid analogs displaying structural similarities to the naturally-occurring glycolipids, such as glycosphingolipids and glycoacylglycerolipids, are capable of eliciting strong immune responses in both herpes simplex virus vaccine and pseudorabies virus vaccine. Some glycolipids have been synthesized from long chain-alkylamines and fatty acids that are linked directly with the sugars through the anomeric carbon atom, to mimic the functions of the naturally occurring lipid residues.

U.S. Pat. No. 4,258,029 granted to Moloney, assigned to the assignee hereof and incorporated herein by reference thereto, teaches that octadecyl tyrosine hydrochloride (OTH) functioned as an adjuvant when complexed with tetanus toxoid and formalin inactivated type I, II and III poliomyelitis virus vaccine. Also, Nixon-George et al. (ref. 24), reported that octadecyl esters of aromatic amino acids complexed with a recombinant hepatitis B surface antigen, enhanced the host immune responses against hepatitis B virus.

Lipidation of synthetic peptides has also been used to increase their immunogenicity. Thus, Wiesmuller (ref. 25) describes a peptide with a sequence homologous to a foot-and-mouth disease viral protein coupled to an adjuvant tripalmityl-S-glyceryl-cysteinylserine, being a synthetic analogue of the N-terminal part of the lipoprotein from Gram negative bacteria. Furthermore, Deres et al. (ref. 26) reported in vivo priming of virus-specific cytotoxic T lymphocytes with synthetic lipopeptide vaccine which comprised of modified synthetic peptides derived from influenza virus nucleoprotein by linkage to a lipopeptide, N-palmityl-S-[2,3-bis(palmitylxy)-(2RS)-propyl-[R]-cysteine (TPC).

## 2. Immunoassays

The about 200 kDa outer membrane protein of the present invention is useful as an immunogen for the generation of anti-200 kDa outer membrane protein antibodies, as an antigen in immunoassays including enzyme-linked immunosorbent assays (ELISA), RIAs and other non-enzyme linked antibody binding assays or procedures known in the art for the detection of anti-bacterial, anti-Moraxella, and anti-200 kDa outer membrane protein antibodies. In ELISA assays, the about 200 kDa outer membrane protein is immobilized onto a selected surface, for example, a surface capable of binding proteins such as the wells of a polystyrene microtiter plate. After washing to remove incompletely adsorbed about 200 kDa outer membrane protein, a nonspecific protein such as a solution of bovine serum albumin (BSA) that is known to be antigenically neutral with regard to the test sample may be bound to the selected surface. This allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific bindings of antisera onto the surface.

The immobilizing surface is then contacted with a sample, such as clinical or biological materials, to be tested in a manner conducive to immune complex (antigen/antibody) formation. This may include diluting the sample with diluents, such as solutions of BSA, bovine gamma globulin (BGG) and/or phosphate buffered saline (PBS)/Tween. The sample is then allowed to incubate for from 2 to 4 hours, at temperatures such as of the order of about 20° to 37° C. Following incubation, the sample-contacted surface is



washed to remove non-immunocomplexed material. The washing procedure may include washing with a solution, such as PBS/Tween or a borate buffer. Following formation of specific immunocomplexes between the test sample and the bound about 200 kDa outer membrane protein, and subsequent washing, the occurrence, and even amount, of immunocomplex formation may be determined by subjecting the immunocomplex to a second antibody having specificity for the first antibody. If the test sample is of human origin, the second antibody is an antibody having specificity for human immunoglobulins and in general IgG. To provide detecting means, the second antibody may have an associated activity such as an enzymatic activity that will generate, for example, a colour development upon incubating with an appropriate chromogenic substrate. Quantification may then be achieved by measuring the degree of colour generation using, for example, a visible spectrophotometer.

#### BIOLOGICAL MATERIALS

Certain plasmids that contain portions of the gene having the open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis* strain 4223 that are described and referred to herein have been deposited with the America Type Culture Collection (ATCC) located at 12301 Parklawn Drive, Rockville, Md., 20852, U.S.A., pursuant to the Budapest Treaty and prior to the filing of this application. The identification of the respective portions of the gene and their molecular size are contained in FIG. 5.

Samples of the deposited plasmids will become available to the public upon grant of a patent based upon this United States patent application. The invention described and claimed herein is not to be limited in scope by plasmids deposited, since the deposited embodiment is intended only as an illustration of the invention. Any equivalent or similar plasmids that encode similar or equivalent antigens as described in this application are within the scope of the invention.

Plasmid	ATCC Designation	Date Deposited
pKS47	97,111	Apr. 7, 1995
pKS5	97,110	Apr. 7, 1995
pKS9	97,114	Apr. 18, 1995

#### EXAMPLES

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific Examples. These Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitations.

Methods of molecular genetics, protein biochemistry, and immunology used but not explicitly described in this disclosure and these Examples are amply reported in the scientific literature and are well within the ability of those skilled in the art.

##### Example 1

This Example illustrates the generation of a non-clumping strain (RH408) of *M. catarrhalis*.

*M. catarrhalis* strain 4223, a clumping strain (a common property of *Moraxella* strains), was inoculated into several flasks containing 20 mL of brain heart infusion (BHI) broth, and the cultures were incubated with shaking (170 rpm) overnight at 37° C. Five mL of each overnight culture were transferred to five individual 1-mL tubes, and were left sitting undisturbed at room temperature for 3 to 8 hours, to allow bacteria to sediment. One hundred  $\mu$ L of the cleared upper phase of each culture medium were used to inoculate 25 mL of BHI broth and cultures were incubated overnight at 37° C., as described above. This passaging was repeated six times, using 25  $\mu$ L of cleared medium to inoculate 25 mL of BHI for each overnight culture. Non-clumping bacterial cultures were identified by measuring the absorbency  $A_{578}$  at intervals over a 3 hour time period, in order to compare the sedimentation rates of the passaged strains to that of the original *M. catarrhalis* strain 4223 culture. Non-clumping mutants, including *M. catarrhalis* RH408, did not aggregate during the three hour time period. On BHI agar plates, strain RH408 had a colony morphology typical for all non-clumping strains. Strain RH408 was previously deposited at the ATCC under the Budapest Treaty on Dec. 13, 1994 with Accession No. 55637.

##### Example 2

This Example illustrates the identification of the above 200 kDa outer membrane protein of *Moraxella catarrhalis*.

*M. catarrhalis* strains 4223, RH408, 5191, 8185, M2, M5, ATCC 25240, 3, 56, 135, 585 were grown in brain heart infusion (BHI) broth. The culture was incubated overnight with aeration at 37° C.

*M. catarrhalis* cells were sonicated and the total protein was determined using the BCA assay system (Pierce, Rockford, Ill.). Ten  $\mu$ g of total protein were mixed with the SDS-PAGE sample buffer containing 0.3M Tris-HCl (pH 8.0), 50% glycerol, 10% SDS, 20%  $\alpha$ -mercaptoethanol and 0.01% bromophenol blue, boiled for 5 minutes and loaded on each lane of SDS-PAGE gel (0.75 mm thick, 7.5% acrylamide). The gels were run at 200 V for 1 hour. Proteins were visualized by staining gels with a solution containing 0.13% Coomassie brilliant blue R, 10% acetic acid and 45% methanol. Excess stain was removed with a destaining solution of 5% ethanol and 7.5% acetic acid.

The various *Moraxella* proteins separated by this procedure are shown in FIGS. 1A and 1B. The *M. catarrhalis* strains tested were as follows:

Lane	Bacterial Strain	Source
FIG. 1A		
1.	Molecular Weight Standards	
2.	<i>E. coli</i>	
3.	No sample	
4.	<i>M. catarrhalis</i> 4223	middle ear fluid
5.	<i>M. catarrhalis</i> RH408	non-clumping variant of 4223
6.	<i>M. catarrhalis</i> 5191	middle ear fluid
7.	<i>M. catarrhalis</i> 8185	nasopharynx
8.	<i>M. catarrhalis</i> M2	sputum
9.	<i>M. catarrhalis</i> M5	sputum
10.	<i>M. catarrhalis</i> 25240	ATCC 25240

FIG. 1B

1.	<i>E. coli</i>
2.	No sample
3.	Molecular Weight Size Markers

-continued

Lane	Bacterial Strain	Source
4.	<i>M. catarrhalis</i> 4223	middle ear fluid
5.	<i>M. catarrhalis</i> RH408	non-clumping variant of 4223
6.	<i>M. catarrhalis</i> 3	sputum
7.	<i>M. catarrhalis</i> 56	sputum
8.	<i>M. catarrhalis</i> 135	middle ear fluid
9.	<i>M. catarrhalis</i> 585	Blood

The about 200 kDa outer membrane protein was clearly seen in all otitis media strains (*M. catarrhalis* 4223, 5191, 135), one isolate from nasopharynx (8185), and in one isolate from sputum (M2). However, the about 200 kDa protein was not detected in three isolates from sputum (3, 56 and M5) and in one strain with unknown origin (ATCC 25240). A very narrow band was found in an isolate from blood of a bacteremia patient (585) and this band was recognized by an anti-200 kDa specific guinea pig serum on an immunoblot. Strain RH408 is a non-clumping spontaneous mutant isolated from strain 4223 (see Example 1) and was found to not express the about 200 kDa protein.

#### Example 3

This Example illustrates the detection of antibodies specific for the about 200 kDa outer membrane protein in a serum obtained from a convalescent patient having recovered from otitis media due to *M. catarrhalis*.

After separation by SDS-PAGE, bacterial proteins were transferred from acrylamide gels to prepared PVDF (polyvinylidene fluoride; Millipore) membranes at a constant voltage of 70 V for 1.5 h in a buffer system consisting of 3 g Tris, 14.4 g glycine and 200 ml methanol per liter at 4° C. Membranes with transferred proteins were blocked with Blocking Reagent (from Boehringer Mannheim) diluted in TBS (0.1M Tris, 0.15M NaCl) at room temperature for 30 min. Blots were exposed to convalescent antiserum diluted 1:500 in Blocking Reagent/TBS with 0.1% Tween 20 for 2 hours at room temperature. This patient had otitis media and the *M. catarrhalis* strain isolated from the patient's ear fluid was *M. catarrhalis* CJ7. Blots were then washed 2 times in Blocking Reagent/TBS with Tween at 15 min per wash. The reporter conjugate, horseradish peroxidase (HRP) conjugated to protein G, was diluted 1:4000 with Blocking Reagent/TBS with Tween and used to immerse the washed membranes for 30 min at room temperature. Blots were washed twice as above, followed by a TBS wash. Bound antibodies were detected using the Lumi-Glo (Kirkegaard and Perry) chemiluminescent detection system as described by the manufacturer. Treated blots were exposed to X-ray film. Antibodies were detected in this convalescent serum that reacted with the about 200 kDa outer membrane protein of *M. catarrhalis* CJ7. These results indicate that the about 200 kDa outer membrane protein is an immunogenic protein of *M. catarrhalis* to which an immune response is elicited during a natural infection by *M. catarrhalis*.

#### Example 4

This Example illustrates the isolation and purification of the about 200 kDa outer membrane protein.

*M. catarrhalis* 4223 cells were harvested by centrifugation at 2,000 rpm for 10 min and frozen. The frozen cells were thawed, resuspended in 20 mM sodium phosphate

buffer (pH 7.2) and sonicated until the cells were disrupted. The frozen-thawed cells were also lysed in 20 mM Tris buffer (pH 8) containing 4% SDS and 0.2 mM EDTA by boiling for 5 min to produce a cell lysate. The cell sonicates and cell lysates were suspended in a SDS-polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer, boiled for 5 min and separated by SDS-PAGE on a gel (1.5 mm thick, 7.5% acrylamide). The estimated position of about 200 kDa protein on the gel was cut out and the protein extracted from the gel by electroelution using the same buffer as the SDS-PAGE running buffer. The isolated about 200 kDa outer membrane protein was shown to be a homogeneous, single band by SDS-PAGE as seen in FIG. 3. The samples analyzed in FIG. 3 are as follows:

Lane	Sample
1.	Molecular Weight Size Markers
2.	Isolated and purified 200 kDa outer membrane protein

The isolated and purified 200 kDa outer membrane protein of *M. catarrhalis* shown in FIG. 3 has a purity of at least 70%. Purified about 200 kDa outer membrane protein preparations of at least 95% could be readily achieved.

When gels were run longer, they showed heterogeneity in the apparent molecular masses of the about 200 kDa outer membrane protein in different strains of *M. catarrhalis* (FIG. 2). In FIG. 2 the strains analyzed were as follows:

Lane	Strain	Source
1.	Molecular Weight Size Markers	
2.	<i>M. catarrhalis</i> H04	middle ear fluid
3.	<i>M. catarrhalis</i> H12	middle ear fluid
4.	<i>M. catarrhalis</i> P034	middle ear fluid
5.	<i>M. catarrhalis</i> P051	middle ear fluid
6.	<i>M. catarrhalis</i> E-07	middle ear fluid
7.	<i>M. catarrhalis</i> E-22	middle ear fluid
8.	<i>M. catarrhalis</i> E-23	middle ear fluid
9.	<i>M. catarrhalis</i> RH 4223	middle ear fluid
10.	<i>M. catarrhalis</i> RH 408	Non-clumping variant of 4223

The strain H12 (lane 3) was a natural isolate from middle ear fluid, but did not produce the about 200 kDa protein.

There may be at least three different sizes of protein in the about 200 kDa range. However, antibodies raised against the about 200 kDa outer membrane protein from one strain of *M. catarrhalis* (4223) did recognize all about 200 kDa proteins tested, present in different strains of *M. catarrhalis*. It is possible, however, that in particular immunogenic compositions, for example, as a vaccine and in particular diagnostic embodiments, that the about 200 kDa outer membrane protein from a variety of *M. catarrhalis* isolates (including immunogenically diverse isolates) may be required.

#### Example 5

This Example illustrates the immunization of guinea pigs with purified about 200 kDa protein from *M. catarrhalis*.

Approximately 30 to 40 µg of the about 200 kDa protein, which was isolated from *M. catarrhalis* strain 4223 by electroelution, were mixed with Freund's complete adjuvant (FCA) and was subcutaneously injected into guinea pigs. After two weeks, the animals were boosted with about the

15

same amount of the about 200 kDa protein in incomplete Freund's adjuvant (IFA). Two weeks later, blood was collected from the guinea pigs and antisera were obtained.

One antiserum was examined on Western blot for its reactivity with the about 200 kDa protein present in 54 different strains of *M. catarrhalis*, which were isolated in different geographical locations throughout the world (Canada, U.S. and Finland) (see Table 1 below). The about 200 kDa protein band was clearly recognized by the antiserum in all strains, in which the presence of the about 200 kDa protein band was detected on SDS-PAGE gels stained with Coomassie Blue. These results indicate that common epitopes of the about 200 kDa protein were conserved in all *M. catarrhalis* strains, which possessed this protein. As stated earlier, this protein is not present in all *M. catarrhalis* strains, but almost all strains, which were isolated from middle ear fluids from otitis media patients, did possess this protein (Table 1).

#### Example 6

This Example illustrates the specific recognition of *M. catarrhalis* strain 4223 with anti-200 kDa protein guinea pig serum by ELISA assay (see Table 2 below).

*M. catarrhalis* strains 4223, RH408 (200 kDa negative mutant) and H-12 were cultured in 60 mL BHI broth overnight. *E. coli* strain BL21 (DE3) was cultured in 60 mL Luria-Bertani (LB) broth overnight. The cultures were split into three tubes and centrifuged. *M. catarrhalis* strain 4223 was centrifuged at 1,500 rpm for 10 minutes, H-12 at 2,000 rpm for 10 minutes, and RH408 and *E. coli* BL21 (DE3) at 3,000 rpm for 15 minutes. The pellet in one tube was suspended in 20 ml of Dulbecco's phosphate buffered saline (D-PBS) and diluted to 1/500 with coating buffer (0.05M carbonate/bicarbonate buffer) pH 9.6. One hundred  $\mu$ L of the bacteria solution was distributed in each well and incubated for 1 hour at room temperature. One hundred  $\mu$ L of 0.2% glutaraldehyde was added to each well and incubated at room temperature for 10 minutes to fix the cells on the well. The wells were washed with PBS containing 0.1% Tween 20 and 0.1% BSA (washing buffer), and then blocked with PBS containing 0.1% BSA for 30 minutes at room temperature. After washing 5 times for 10 seconds with the washing buffer, serial dilutions of guinea pig antiserum with the washing buffer were added to wells and incubated at room temperature for 60 minutes. After washing, goat anti-guinea pig IgG conjugated with horseradish peroxidase was added to each well at the dilution of 1/20,000. After incubation at room temperature for 60 minutes, the wells were washed and then color reaction was developed using 3,3',5,5'-tetramethylbenzidine (TMB) and hydrogen peroxide.

The ELISA plate wells were also coated with sonicates containing 10  $\mu$ g/mL of total proteins in the coating buffer, blocked without the fixation process and then assayed as described above.

The results shown in Table 2 indicate that the 200 kDa outer membrane protein specific guinea pig antiserum specifically recognizes strains of *M. catarrhalis* which produce the about 200 kDa protein. The ability of the antiserum to recognize whole cells indicates that the protein is present on the surface of the bacterial cells.

#### Example 7

This Example describes the determination of an internal amino acid sequence of the 200 kDa outer membrane protein.

The about 200 kDa outer membrane protein was isolated from *M. catarrhalis* 4223 as described above. The protein

16

was subjected to CNBr degradation, the proteolytic digests subjected to SDS-PAGE and transferred onto PVDF membrane. A peptide band migrating at a position corresponding to approximately 40 kDa was cut out from the membrane and its N-terminal amino acid sequence was determined. In another experiment, the CNBr degradation products of the about 200 kDa protein were subjected to a direct determination of N-terminal amino acid sequencing without separating by SDS-PAGE. Both analyses gave an identical, N-terminal sequence of 20 amino acids with one unidentified amino acid at the 17th position. The internal sequence of the 200 kDa outer membrane protein was:

NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-X-Gln-Gly-Ile (SEQ ID No: 2).

#### Example 8

This Example describes the immunization of guinea pigs with a peptide corresponding to an internal fragment of the about 200 kDa outer membrane protein and the analysis of the antiserum generated.

Based upon the determination of the amino acid sequence of an internal fragment of the about 200 kDa outer membrane protein, a 16 amino acid long peptide of sequence:

NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys (SEQ ID No: 3)

was synthesized by standard procedure. This 16-mer peptide was conjugated to KLH using Impject Maleimide Activated KLH (Pierce, Rockford, Ill.) and approximately 500  $\mu$ g of the conjugate was injected into guinea pigs using the same immunization and boosting schedule as described above. The guinea pig serum raised against the 16-mer internal amino acid sequence (SEQ ID No: 3) was examined by immunoblot analysis and found to specifically recognize 200 kDa outer membrane protein in cell sonicates of *M. catarrhalis* 4223. The results are shown in FIG. 4 and indicate that the anti-peptide guinea pig antiserum specifically recognizes the 200 kDa protein of *M. catarrhalis* 4223. The samples analyzed in FIG. 4 were as follows:

Lane	Sample	Antiserum
1.	Molecular Weight Markers	
2.	Purified 200 kDa outer membrane protein	Anti-200 kDa protein
3.	<i>M. catarrhalis</i> cell sonicate	Anti-peptide 1:5000
4.	<i>M. catarrhalis</i> cell sonicate	Anti-peptide 1:1000
5.	<i>M. catarrhalis</i> cell sonicate	Anti-peptide 1:500
6.	<i>M. catarrhalis</i> cell sonicate	Pre-immune serum

The results obtained confirm that the peptide corresponding to SEQ ID Nos: 2 and 3 are derived from the 200 kDa outer membrane protein.

#### Example 9

This Example describes the preparation of a *M. catarrhalis* library.

Chromosomal DNA was isolated as follows:

*M. catarrhalis* cell pellet was resuspended in 20 mL Tris-EDTA (TE) buffer, pH 7.5. Pronase (final concentration 500  $\mu$ g/mL) and SDS (final concentration 1%) were added and the suspension was incubated in a 37° C. water bath for 2 hours. DNA was isolated by sequential extractions with phenol (1 time), phenol-chloroform (1:1, 2 times), and chloroform-isoamyl alcohol (24:1, 1 time). Extracted DNA was dialyzed against 1M NaCl at 4° C. for 4 hours. This was

17

followed by dialysis against TE buffer, pH 7.5, at 4° C. for 48 hours (3 buffer changes). DNA was ethanol precipitated from the dialysate. Large size DNA was collected by spooling on a glass rod. Air dried DNA was dissolved in 3 mL water. Small scale Sau3A (New England BioLabs) restriction digests of chromosomal DNA (final volume 10  $\mu$ L) were done to establish conditions required to obtain maximal amounts of chromosomal DNA with a size range of 15–23 kb. Large scale digests were prepared once the optimal digestion conditions were defined. The large scale digests consisted of 50  $\mu$ L chromosomal DNA (290  $\mu$ g/mL), 33  $\mu$ L water, 10  $\mu$ L Sau3A buffer (New England BioLabs), 1  $\mu$ L BSA (10 mg/mL, New England BioLabs) and 6.3  $\mu$ L Sau3A (0.04 U/ $\mu$ L), and were incubated at 37° C. for 15 minutes. Reactions were stopped with the addition of 10  $\mu$ L 10X loading buffer (100 mM Tris-HCl pH 8, 10 mM EDTA, 0.1% bromophenol blue, 50% glycerol). Digested DNA was applied to 0.5% agarose gels (prepared in Tris-acetate-EDTA (TAE)) and separated according to size at 50 V for 6 hours. The region of the gel encompassing DNA of size 15–23 kb was cut from the gel and placed in dialysis tubing (BRL) with 3 mL TAE. DNA was electroeluted from the gel slice overnight at a field strength of 1 V/cm. Electroeluted DNA in TAE was extracted once with phenol, once with phenol-chloroform (1:1), and finally precipitated with ethanol. The dried DNA pellet was dissolved in 5  $\mu$ L water. Size-fractionated chromosomal DNA was ligated with BamHI cut EMBL3 arms (Promega) using T4 DNA ligase in a final volume of 9  $\mu$ L. The entire ligation reaction was packaged into phage  $\lambda$  using a commercially purchased packaging kit following the manufacturer's (Amersham) protocol.

The packaged DNA library was amplified on solid medium. This was accomplished by incubating 0.1 mL *E. coli* strain NM539 plating cells (cells suspended in 10 mM MgSO<sub>4</sub>) with 15–25  $\mu$ L of the packaged DNA library at 37° C. for 15 minutes. Bacteria with adsorbed phage were plated onto BBL plates (10 g BBL trypticase peptone, 5 g NaCl and 15 g agar per liter) using 3 mL of BBL top-agarose (same as BBL plates except agar replaced with 0.6% agarose) and plates were incubated overnight at 37° C. Phage were eluted from the top-agarose by adding 3 mL SM buffer (50 mM Tris-HCl, pH 7.5, 8 mM MgSO<sub>4</sub>, 100 mM NaCl, 0.01% gelatin) to the plates and leaving them at 4° C. for 7 hours. SM buffer containing phage was collected from the plates, transferred to a screwcap tube and stored at 4° C. over chloroform.

#### Example 10

This Example describes the cloning on a gene encoding the *M. catarrhalis* 200 kDa outer membrane protein.

The *M. catarrhalis* genomic library in phage lambda EMBL3 was screened using an anti-200 kDa protein guinea pig antiserum. A lambda phage clone 8II (FIG. 5), which expressed as about 200 kDa protein, was confirmed by immunoblotting of the phage lysate using the about 200 kDa outer membrane-specific antiserum.

Plate lysate cultures of this recombinant phage were prepared. The DNA was extracted from the plate lysates using a Wizard Lambda Preps DNA Purification System (Promega Corp., Madison, Wis.) according to the manufacturer's instructions. This phage clone carried a DNA insert of about 16 kb in size (the restriction map for which is in FIG. 5). The phage DNA was digested with a mixture of two restriction enzymes, Sall and XhoI, and separated by agarose gel electrophoresis. Two DNA bands, approximately 5 kb and 11 kb in size, respectively, were cut out from the gel

18

and extracted using a GeneClean kit (BIO 101 Inc., LaJolla, Calif.) according to a manufacturer's instruction.

The smaller 5 kb fragment was ligated into a plasmid vector, pBluescript II SK +/- (Stratagene Cloning Systems, LaJolla, Calif.), which was previously digested with Sall and XhoI, to produce plasmid pKS5. The larger 11 kb fragment was ligated into a plasmid vector, pSP72 (Promega Corp., Madison, Wis.), to produce plasmid pKS9. Both ligated plasmids were used to transform *E. coli*, strain DH5 $\alpha$ .

The lambda phage DNA was also digested with a mixture of XhoI and KpnI and the approximately 1.2 kb fragment was isolated after agarose gel separation as described above. This 1.2 kb fragment was ligated into a plasmid vector, pGEM-7Zf(+) (Promega Corp., Madison, Wis.), to produce plasmid pKS47. Restriction maps of the lambda and plasmid clones are shown in FIG. 5.

#### Example 11

This Example describes the sequencing of the gene having an open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis*.

The gene encoding the about 200 kDa outer membrane protein was sequenced by an Applied Biosystems sequencer. The one strand of the insert in the plasmid pKS5, was sequenced after construction of a nested set of deletions using a Erase-a-Base system (Promega Corp., Madison, Wis.). The plasmid pKS5 was first digested with XhoI and KpnI, treated with exonuclease III to generate a nested set of deletions in the insert and then recircularized according to the manufacturer's instructions. *E. coli* DH5 $\alpha$  was transformed with a series of plasmids with deletions generated in this way. Plasmids were isolated from the transformants using a Qiagen midi plasmid isolation kit (Qiagen) and the size of plasmids examined by agarose gel electrophoresis after restriction enzyme digestion. The inserts of the plasmids with deletions were sequenced using a bacteriophage T7 promoter sequence as a primer.

Based upon the sequence, nucleotide primers were synthesized. Using the synthetic nucleotide primers, sequence gaps, which were not sequenced by the Erase-a Base system, were determined.

The sequence of the insert in plasmid pKS47 was determined from both ends using synthetic nucleotide primers. The nucleotide sequence of the gene has an open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis* as shown in FIG. 6 (SEQ ID No: 1). This sequence included a nucleotide sequence:

5'-AATGTCAAATCAGTCATTAACAAAGAACAAG  
TAAATGATGCCAATAAAAAGCAAGGCATC-3'  
(SEQ ID No: 4) which encodes the internal amino acid sequence of the about 200 kDa outer membrane protein (SEQ ID No: 2) determined above. This result confirms that the cloned gene has an open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis*.

#### SUMMARY OF THE DISCLOSURE

In summary of the disclosure, the present invention provides an isolated and purified outer membrane protein of a Moraxella strain, particularly *M. catarrhalis*, having a molecular weight of about 200 kDa as well as isolated and purified DNA molecules encoding the outer membrane protein. The invention also provides peptides corresponding to portions of the outer membrane protein. The protein, DNA

sequences, recombinant proteins derived therefrom and peptides are useful for diagnosis, immunization and the generation of diagnostic and immunological reagents. Modifications are possible within the scope of this invention.

TABLE I

Presence of about 200 kDa outer membrane protein in various isolates of <i>Moraxella catarrhalis</i>		
Type of Clinical Isolate	Number of isolates Examined	Number of isolates <sup>1</sup> containing the 200 kDa outer membrane protein
Otitis Media	37	36
Sputum/Expectoration/ Bronchial Secretion	13	6
Blood	2	2
Nasopharynx	1	1
Unknown	1	0

<sup>1</sup>The presence of the about 200 kDa outer membrane protein was determined by immunoblot analysis using a monospecific guinea pig anti-200 kDa protein antiserum.

TABLE II

Detection of about 200 kDa outer membrane protein of <i>M. catarrhalis</i> by the monospecific anti-200 kDa outer membrane guinea pig antiserum		
Strain	Sample	Reciprocal Reactive Titre
4223	Whole cells not fixed	800
RH408	Whole cells not fixed	<200
H12	Whole cells not fixed	<200
<i>E. coli</i> BL21	Whole cells not fixed	<200
4223		3200
RH408		200
H12		<200
<i>E. coli</i> BL21		<200
4223	Sonicate	12,800
RH408	Sonicate	800
H12	Sonicate	800
<i>E. coli</i> BL21	Sonicate	200

## REFERENCES

- Van Hare, G. F., P. A. Shurin, C. D. Marchant, N. A. Cartelli, C. E. Johnson, D. Fulton, S. Carlin, and C. H. Kim. Acute otitis media caused by *Branhamella catarrhalis*: biology and therapy. *Rev. Infect. Dis.* 9:16-27.
- Chapman, A. J., D. M. Musher, S. Jonsson, J. E. Clarridge, and R. J. Wallace. 1985. Development of bactericidal antibody during *Branhamella catarrhalis* infection. *J. Infect. Dis.* 151:878-882.
- Hager, H., A. Verghese, S. Alvarez, and S. L. Berk. 1987. *Branhamella catarrhalis* respiratory infections. *Rev. Infect. Dis.* 9:1140-1149.
- McLeod, D. T., F. Ahmad, M. J. Croughan, and M. A. Calder. 1986. Bronchopulmonary infection due to *M. catarrhalis*. Clinical features and therapeutic response. *Drugs* 31(Suppl. 3):109-112.
- Nicotra, B., M. Rivera, J. I. Luman, and R. J. Wallace. 1986. *Branhamella catarrhalis* as a lower respiratory tract pathogen in patients with chronic lung disease. *Arch. Intern. Med.* 146:890-893.
- Ninane, G., J. Joly, and M. Kraytman. 1978. Bronchopulmonary infection due to *Branhamella catarrhalis* 11 cases assessed by transtracheal puncture. *Br. Med. J.* 1:276-278.
- Srinivasan, G., M. J. Raff, W. C. Templeton, S. J. Givens, R. C. Graves, and J. C. Mel. 1981. *Branhamella catarrhalis* pneumonia. Report of two cases and review of the literature. *Am. Rev. Respir. Dis.* 123:553-555.
- West, M., S. L. Berk, and J. K. Smith. 1982. *Branhamella catarrhalis* pneumonia. *South. Med. J.* 75:1021-1023.
- Brorson, J.-E., A. Axelsson, and S. E. Holm. 1976. Studies on *Branhamella catarrhalis* (*Neisseria catarrhalis*) with special reference to maxillary sinusitis. *Scan. J. Infect. Dis.* 8:151-155.
- Evans, F. O., Jr., J. B. Sydnor, W. E. C. Moore, G. R. Moore, J. L. Manwaring, A. H. Brill, R. T. Jackson, S. Hanna, J. S. Skaar, L. V. Holdeman, G. S. Fitz-Hugh, M. A. Sande, and J. M. Gwaltney, Jr. 1975. Sinusitis of the maxillary antrum. *N. Engl. J. Med.* 293:735-739.
- Tinkelman, D. G., and H. J. Silk. 1989. Clinical and bacteriologic features of chronic sinusitis in children. *Am. J. Dis. Child.* 143:938-942.
- Wald, E. R., C. Byers, N. Guerra, M. Casselbrant, and D. Beste. 1989. Subacute sinusitis in children. *J. Pediatr.* 115:28-32.
- Wald, E. R., G. J. Milmo, A. Bowen, J. Ledesma-Medina, N. Salamon, and C. D. Bluestone. 1981. Acute maxillary sinusitis in children. *N. Engl. J. Med.* 304:749-754.
- Christensen, J. J., and B. Bruun. 1985. Bacteremia caused by a beta-lactamase producing strain of *Branhamella catarrhalis*. *Acta. Pathol. Microbiol. Immunol. Scand. Sect. B* 93:273-275.
- Craig, D. B., and P. A. Wehrle. 1983. *Branhamella catarrhalis* septic arthritis. *J. Rheumatol.* 10:985-986.
- Gray, L. D., R. E. Van Scoy, J. P. Anhalt, and P. K. W. Yu. 1989. Wound infection caused by *Branhamella catarrhalis*. *J. Clin. Microbiol.* 27:818-820.
- Guthrie, R., K. Bakenhaster, R. Nelson, and R. Woskobnick. 1988. *Branhamella catarrhalis* sepsis: a case report and review of the literature. *J. Infect. Dis.* 158:907-908.
- Hiroshi, S., E. J. Anaissie, N. Khardori, and G. P. Bodey. 1988. *Branhamella catarrhalis* septicemia in patients with leukemia. *Cancer* 61:2315-2317.
- O'Neill, J. H., and P. W. Mathieson. 1987. Meningitis due to *Branhamella catarrhalis*. *Aust. N.Z. J. Med.* 17:241-242.
- Murphy, T. F. 1989. The surface of *Branhamella catarrhalis*: a systematic approach to the surface antigens of an emerging pathogen. *Pediatr. Infect. Dis. J.* 8:S75-S77.
- Klingman, K. L., and T. F. Murphy. 1994. Purification and characterization of a high-molecular-weight outer membrane protein of *Moraxella* (*Branhamella*) *catarrhalis*. *Infect. Immun.* 62:1150-1155.
- Helminen, M. E., I. Maciver, J. L. Latimer, J. Klesney-Tait, L. D. Cope, M. Paris, G. H. McCracken, Jr., and E. J. Hansen. 1994. A large, antigenically conserved protein on the surface of *Moraxella catarrhalis* is a target for protective antibodies. *J. Infect. Dis.* 170:867-872.
- Panezutti H., O. James, E. J. Hanson, Y. Choi, R. E. Harkness, M. H. Klein and P. Chong. 1993. Identification of surface-exposed B-cell epitopes recognized by *Haemophilus influenzae* type b P1 specific monoclonal antibodies. *Infect. Immun.* 61: 1867-1872.
- Nixon-George et al. (1990), *J. Immunology* 144:4798-4802.
- Wiesmuller (1989), *Vaccine* 8:29-33.
- Deres et al. (1989), *Nature* 342:561.
- Lockhoff, O. Glycolipids as Immunomodulators: Synthesis and Properties. 1991. *Chem. Int. Ed. Engl.* 30:1611-1620.

## SEQUENCE LISTING

&lt;160&gt; NUMBER OF SEQ ID NOS: 4

&lt;210&gt; SEQ ID NO 1

&lt;211&gt; LENGTH: 6975

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Moraxella catarrhalis*

&lt;400&gt; SEQUENCE: 1

```

ccatggatat gggcagggtg gctgcctgc cgtatgatgg cgtacacacc ccatttgcgc 60
catatctgta cgatttgaca tgtgatatga tttaacatgt gacatgattt aacattgttt 120
aatactgttg ccatcattac cataatttag taacgcattt agtaacgcat ttgtaaaaat 180
cattgcgcgc ctttatgtgt atcatatgaa tagaatatta tgattgtatc tgattattgt 240
atcagaatgg tgatgctata tgatgatgcc tacgagttga tttgggttaa tcactctatg 300
atttgatata ttttgaaact aatctattga cttaaatcac catatggtta taatttagca 360
taatggtagg cttttttaa aaatcacatc gcaatattgt tctactgtta ctaccatgct 420
tgaatgacga tcccaatcac cagattcatt caagtgatgt gttgtatac gcaccattta 480
ccctaattat ttcaatcaaa tgcctatgtc agcatgtatc atttttttaa ggtaaaccac 540
catgaatcac atctataaag tcatctttaa caaagccaca ggcacattta tggcagtggc 600
agagtacgcc aaatccaca gcacgggggg ggggtagctg tgctacaggg caagttggca 660
gtgtatgcac tctgagcttt gccgtattg ccgcgctcgc tgcctcgtg atcggtgcaa 720
cgctcagtg gctgcttat gctcaaaaaa aagataccaa acatcgcga attggtgaac 780
aaaaccagcc aagacgctca ggcactgcca aggcggacgg tgatcgagcc attgctattg 840
gtgaaaatgc taacgcacag ggcgggcaag ccacgcgat cggtagtagt aataaaactg 900
tcaatggaag cagtttgat aagataggta ccgatgtac gggcaagag tccatcgcca 960
tcgggtgtga tgtaaaggct agtgggtgat cctcgattgc catcggtagt gatgacttac 1020
atttgcttga tcagcatggt aatcctaacc atccgaaagg tactctgatt aacgatctta 1080
ttaacggcca tgcagtatta aaagaaatc gaagctcaaa ggataatgat gtaaaatata 1140
gacgcacaac cgcaagcgga cagccagta ctgcagtggg agccatgtca tatgcacagg 1200
gtcatttttc caacgccttt ggtacacggg caacagctaa aagtcctat tccttggcag 1260
tgggtcttgc cgccacagcc gagggccaat ctacaatgc tattgttct gatgcaacat 1320
ctagctcgtt gggagcgata gcccttggg caggtactcg tgctcagcta cagggcagta 1380
ttgccctagg tcaaggctt gttgtcactc agagtataa taattctaga cgggctata 1440
caccaaatcc ccaggcacta gacccaagt ttcaagccac caataatcg aagcggggtc 1500
cactttccat tggtagtaac tctatcaaac gtaaaatcat caatgtcggg gcaggtgtta 1560
ataaaaccga tgcgggcaat gtggcacagc tagaagcggg ggtgaagtgg gctaaggagc 1620
gtagaattac ttttcagggt gatgataaca gtactgacgt aaaaataggt ttggataata 1680
ctttaactat taaagggtgt gcagagacca acgcattaac cgataataat atcgggtgtg 1740
taaaagaggg tgataatagt ggtctgaaag ttaacttgc taaacttta aacaatctta 1800
ctgaggtgaa tacaactaca ttaaatgcca caaccacagt taaggtaggt agtagtagta 1860
gtactacagc tgaattattg aagtagatt taacctttac ccagcccaat acaggcagtc 1920
aaagcacaag caaaaccgtc tatggcgtaa atgggggtgaa gtttactaat aatgcagaaa 1980
caacagcagc aatcggcact actcgtatta ccagagataa aattggcttt gtcgagatg 2040

```

-continued

gtgatgttga tgaanaacaa gcaccatatt tggataaaaa acaacttaaa gtgggtagtg 2100  
 ttgcaattac catagacaat ggcattgatg caggtataaa aaagatcagt aatcttgcca 2160  
 aaggtagcag tgctaacgat gcggttacca tcgaacagct caaagccgcc aagcctactt 2220  
 taaacgcagg cgctggcacc agtgtcacac ctactgaaat atcagttgat gctaagagt 2280  
 gcaatgttac cgccccaact tacaacattg gcgtgaaaac caccgagctt aacagtgatg 2340  
 gcactagtga taaatttagt gttaaaggga gtggtacgaa caatagctta gttaccgccg 2400  
 aacatttgcc aagctatcta aatgaagta atcgaacggc tgacagtgtc ctacaaagct 2460  
 ttaccggtta agaagaagac gatgatgacg ccaacgctat caccgtggct aaagatacga 2520  
 caaaaaatgc cggcgagcgc agcatcttaa aactcaaagg taaaaacggg ctaacggttg 2580  
 ctacaaaaaa agatggtaac gttacccttg ggcttagcca agatagcggg ctgaccattg 2640  
 gcaaaagcac cctaaacaac gatggcttga ctgttaaaga taccaacgaa caaatccaag 2700  
 tcggtgctaa tggcattaaa ttactaatg tgaatgtag taatccaggt actggcattg 2760  
 caaataccgc tcgcattacc agagataaaa ttggctttgc tggttctgat ggtgcagttg 2820  
 atacaaacaa accttatctt gatcaagaca agctacaagt tggcaatgtt aagattacca 2880  
 aactggcat taacgcaggt ggtaaagcca tcacagggct gtcccaaca ctgcctagca 2940  
 ttgcccagca aagtagccgc aacatagaac tgggcaatc aatccaagac aaagacaaat 3000  
 ccaacgctgc cagcattaat gatataata atacaggctt taacctaaaa aataataaca 3060  
 accccattga ctttgtctcc acttatgaca ttgttgactt tgccaatggc aatgccacca 3120  
 ccgccacagt aaccatgat accgctaaca aaaccagtaa agtggatat gatgtgaatg 3180  
 tggatgatac aaccattcat ctaacaggca ctgatgaca taaaaaactt ggcgtcaaaa 3240  
 ccaccaaact gaacaaaaca agtgcataat gtaatacagc aactaacttt aatgttaact 3300  
 ctatgtatga agatgccctt gtaaacgcca aagacatcgc cgaaaatcta aacaccctag 3360  
 ccaaggaaat tcacaccacc aaaggcacag cagacaccgc cctacaaacc ttaccgtta 3420  
 aaaaaggtaga tgaanaaat aatgctgatg acgccaacgc catcaccgtg ggtcaaaaaga 3480  
 acgcaataaa tcaagtcaac accctaacac tcaaaaggta aaacggctctt aatattaaaa 3540  
 ccgacaaaaa tggtagcgtt acctttggca ttaacaccac aagcggctctt aaagccggca 3600  
 aaagcaccct aaacgcaggt ggcttgtcta ttaaaaaccc cactggtagc gaacaaatcc 3660  
 aagtcggtgc tgatggcgtg aagtttgcca aggttaataa taatgggtgt gtaggtgctg 3720  
 gcattgatgg cacaactcgc attaccagag atgaattgg ctttactggg actaatggct 3780  
 cacttgataa aagcaaaacc cacctaagca aagacggcat taacgcaggt ggtaaaaaga 3840  
 ttaccaacat tcaatcaggt gagattgccc aaacagcca tgatgctgtg acaggcggca 3900  
 agatttatga tttaaaaacc gaacttgaaa acaaaatcag cagtactgcc aaacagcac 3960  
 aaaactcatt acacgaattc tcagtagcag atgaacaagg taataacttt acggttagta 4020  
 acccttactc cagttatgac acctcaaga cctctgatgt catcaccttt gcaggtgaaa 4080  
 acggcattac caccaaggta aataaagggt tgggtcgtgt gggcattgac caaaccaaag 4140  
 gcttaaccac gcctaagctg accgtgggta ataataatgg caaaggcatt gtcattgaca 4200  
 gccaaaatgg tcaaaatacc atcacaggac taagcaacac tctagcta atgttaccatg 4260  
 ataaaggtag cgtacgcacc acagaacag gcaatataat caaagacgaa gacaaaaccc 4320  
 gtgccgccag cattgttgat gtgctaagcg caggctttaa cttgcaaggc aatggggaag 4380

-continued

cggttgactt	tgtctccact	tatgacaccg	tcaactttgc	cgatggcaat	gccaccaccg	4440
ctaaggtgac	ctatgatgac	acaagcaaaa	ccagtaaaagt	ggtctatgat	gtcaatgtgg	4500
atgatacaac	cattgaagtt	aaagataaaa	aacttggcgt	aaaaaccacc	acattgacca	4560
gtactggcac	aggtgctaata	aaatttgccc	taagcaatca	agctactggc	gatgcgcttg	4620
tcaaggccag	tgatatcggt	gctcatctaa	acaccttata	tggcgacatc	caaactgcca	4680
aaggggcaag	ccaagcgaac	aactcagcag	gctatgtgga	tgctgatggc	aataaggtca	4740
tctatgacag	taccgataac	aagtactata	aagccaaaaa	tgatggcaca	gttgataaaa	4800
ccaaagaagt	tgccaaagac	aaactggctg	cccaagccca	aaccccgat	ggcacattgg	4860
ctcaaatgaa	tgtaaatca	gtcattaaca	aagaacaagt	aatgatgcc	aataaaaagc	4920
aaggcatcaa	tgaagacaac	gcctttgtta	aaggacttga	aaaagccgct	tctgataaca	4980
aaaccaaaaa	cgccgcagta	actgtgggtg	atttaaatgc	cgttgcccac	acaccgctga	5040
cctttgcagg	ggatacaggc	acaacggcta	aaaaactggg	cgagactttg	accatcaaa	5100
gtgggcaaac	agacaccaat	aagctaaccg	ataataacat	cggtgtggta	gcaggctactg	5160
atggtctcac	tgtaaaactt	gccaagacc	taaccaatct	taacagcgtt	aatgcagggtg	5220
gcacaaaaat	tgatgacaaa	ggcgtgtctt	ttgtagactc	aagcgggtcaa	gccaaagcaa	5280
acacccctgt	gctaagtgcc	aatgggctgg	acctgggtgg	caaggtcatc	agtaatgtgg	5340
gcaaaggcac	aaaagatacc	gaccgctgcc	aatgtacaac	agttaaacga	agtacgcac	5400
ttgttgggtc	ttgtaatagc	tggtaatgat	aacgtgacg	gcaatcaggt	aaacattgcc	5460
gacatcaaaa	aagacccaaa	ttcaggttca	tcacttaacc	gcactgtcat	caaagcaggc	5520
acggtacttg	gcggtaaagg	taataacgat	accgaaaaac	ttgccactgg	tggtatacaa	5580
gtgggctgtg	ataaagacgg	caacgctaac	ggcgatttaa	gcaatgtttg	ggtcaaaacc	5640
caaaaagatg	gcagcaaaaa	agccctgctc	gccacttata	acgccgcagg	tcagaccaac	5700
tatttgacca	acaacccccc	agaagccatt	gacagaataa	atgaacaagg	tatccgcttc	5760
ttccatgtca	acgatggcaa	tcaagagcct	gtggtacaag	ggcgtaacgg	cattgactca	5820
agtgcctcag	gcaagcactc	agtggcgata	ggtttccagg	ccaaggcaga	tggtgaagcc	5880
gccgttgcca	taggcagaca	aaccaagca	ggcaaccaat	ccatcgccat	cggtgataac	5940
gcacaagcca	cgggcatcat	atccatcgcc	atcggtacag	gcaatgtggt	agcaggtaag	6000
cactctggtg	ccatcgggca	cccaagcact	gttaaggctg	ataacagtta	cagtgtgggt	6060
aataacaacc	agtttaaccg	tgccactcaa	accgatgtct	ttggtgtggg	caataacatc	6120
accgtgaccg	aaagtaactc	ggttgcccta	ggttcaaaact	ctgccatcag	tcagggcaca	6180
cacgcaggca	cacaagccaa	aaaatctgac	ggcacagcag	gtacaaccac	cacagcaggt	6240
gcaaccggta	cgggttaaagg	ctttgctgga	caaacggcgg	ttggtgcggt	ctccgtgggt	6300
gcctcagggt	ctgaacgccg	tatccaaaat	gtggcagcag	gtgaggtcag	tgccaccagc	6360
accgatgcgg	tcaatggtag	ccagttgtac	aaagccaccc	aaagcattgc	caacgcaacc	6420
aatgagcttg	accatcgtat	ccaccaaaac	gaaaataagg	ccaatgcagg	gatttcacat	6480
gcgatggcga	tggcgtccat	gccacaagcc	tacattcctg	gcagatccat	ggttaccggg	6540
ggtattgcca	cccacaacgg	tcaaggtgcg	gtggcagtg	gactgtcgaa	gctgtcggat	6600
aatggtcaat	gggtatttaa	aatcaatggt	tcagccgata	cccaaggcca	tgtagggcgg	6660
gcagttgggt	caggttttca	cttttaagcc	ataaatcgca	agattttact	taaaaatcaa	6720
tctcaccata	gttgataaaa	aacagcatca	gcacagtc	tattactgat	gctgatgttt	6780



-continued

```

tttatcactt aaaccatttt accgctcaag tgattctctt tcaccatgac caaatcgcca 6840
ttgatcatag gtaaaccttat tgagtaaatt ttatcaatgt agttgttaga tatggttaaa 6900
attgtgccat tgacccaaaa atgaccgatt tatcccgaaa atttctgatt atgatccgtt 6960
gacctgcagg tcgac 6975

```

```

<210> SEQ ID NO 2
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Moraxella catarrhalis
<220> FEATURE:
<221> NAME/KEY: UNSURE
<222> LOCATION: (17)

```

```

<400> SEQUENCE: 2

```

```

Asn Val Lys Ser Val Ile Asn Lys Glu Gln Val Asn Asp Ala Asn Lys
 1             5             10             15
Xaa Gln Gly Ile
                20

```

```

<210> SEQ ID NO 3
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Moraxella catarrhalis

```

```

<400> SEQUENCE: 3

```

```

Asn Val Lys Ser Val Ile Asn Lys Glu Gln Val Asn Asp Ala Asn Lys
 1             5             10             15

```

```

<210> SEQ ID NO 4
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Moraxella catarrhalis

```

```

<400> SEQUENCE: 4

```

```

aatgtcaaat cagtcattaa caaagaacaa gtaaatgatg ccaataaaaa gcaaggcacc 60

```

What we claim is:

1. An isolated and purified outer membrane protein of a *Moraxella catarrhalis* strain having an apparent molecular mass of about 200 kDa, as determined by SDS-PAGE. 45
2. The protein of claim 1 wherein the strain is *Moraxella catarrhalis* 4223 or RH408.
3. The protein of claim 1 containing the amino acid sequence NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID NO:2) 50 for the *Moraxella catarrhalis* strain 4223.
4. The protein of claim 1 which is at least about 70 wt % pure.
5. The protein of claim 4 which is at least about 95 wt % pure.
6. The protein of claim 1 in the form of an aqueous 55 solution thereof.
7. The protein of claim 1 recognizable by an antibody preparation specific for a peptide consisting of the amino acid sequence of NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID NO:2). 60
8. A recombinant outer membrane protein of a *Moraxella catarrhalis* strain having an apparent molecular mass of about 200 kDa as determined by SDS-PAGE producible by a transformed host containing an expression vector comprising a nucleic acid sequence which is: 65

(A) a purified and isolated nucleic acid molecule encoding an outer membrane protein of a strain of *Moraxella*

*catarrhalis* having an molecular mass of about 200 kDa, as determined by SDS-PAGE, or

(B) a purified and isolated nucleic acid molecule having a sequence selected from the group consisting of:

(a) the nucleotide sequence set out in FIG. 6 (SEQ ID NO:1) or the complementary sequence thereto,

(b) a nucleotide sequence encoding a 200 kDa protein of a strain of *Moraxella catarrhalis* and containing the amino acid sequence NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID NO:2); and

(c) a nucleotide sequence encoding a 200 kDa portion of a strain of *Moraxella catarrhalis* and which hybridizes under stringent conditions to any one of the sequences defined in (a) or (b); and

expression means operatively coupled to the nucleic acid molecule for expression by the host of the 200 kDa outer membrane protein of *Moraxella catarrhalis*.

9. A peptide consisting of the amino acid sequence NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No:2) or NH<sub>2</sub>-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys (SEQ ID No:3) for the *Moraxella catarrhalis* 4223 strain.

\* \* \* \* \*